



Guidelines for the Use of Antiretroviral Agents in Pediatric HIV Infection

Downloaded from <http://aidsinfo.nih.gov/guidelines> on 12/21/2016

Visit the AIDSinfo website to access the most up-to-date guideline.

Register for e-mail notification of guideline updates at <http://aidsinfo.nih.gov/e-news>.

Management of Medication Toxicity or Intolerance (Last updated

March 1, 2016; last reviewed March 1, 2016)

Panel's Recommendations

- In children who have severe or life-threatening toxicity (e.g., a hypersensitivity reaction), all antiretroviral (ARV) drugs should be stopped immediately (AIII). Once symptoms of toxicity have resolved, antiretroviral therapy should be resumed with substitution of a different ARV drug or drugs for the offending agent(s) (AII*).
- When modifying therapy because of toxicity or intolerance to a specific drug in children with virologic suppression, changing one drug in a multidrug regimen is permissible; if possible, an agent with a different toxicity and side-effect profile should be chosen (AI*).
- The toxicity and the medication presumed responsible should be documented in the medical record and the caregiver and patient advised of the drug-related toxicity (AIII).
- Dose reduction is not a recommended option for management of ARV toxicity, except for those few ARV drugs (e.g., efavirenz) for which a therapeutic range of plasma concentrations detected by therapeutic drug monitoring correlates with toxicity (AII*).

Rating of Recommendations: A = Strong; B = Moderate; C = Optional

Rating of Evidence: I = One or more randomized trials in children[†] with clinical outcomes and/or validated endpoints; I* = One or more randomized trials in adults with clinical outcomes and/or validated laboratory endpoints with accompanying data in children[†] from one or more well-designed, nonrandomized trials or observational cohort studies with long-term clinical outcomes; II = One or more well-designed, nonrandomized trials or observational cohort studies in children[†] with long-term outcomes; II* = One or more well-designed, nonrandomized trials or observational studies in adults with long-term clinical outcomes with accompanying data in children[†] from one or more similar nonrandomized trials or cohort studies with clinical outcome data; III = Expert opinion

[†] Studies that include children or children/adolescents but not studies limited to post-pubertal adolescents

Medication Toxicity or Intolerance

The goals of antiretroviral therapy (ART) include achieving and maintaining viral suppression and improving immune function, with a regimen that is not only effective but also as tolerable and safe as possible. This requires consideration of the toxicity potential of a ART regimen, as well as the individual child's underlying conditions, concomitant medications, and prior history of drug intolerance or viral resistance.

Adverse effects (AEs) have been reported with use of all antiretroviral (ARV) drugs, and are among the most common reasons for switching or discontinuing therapy, and for medication nonadherence. However, rates of treatment-limiting AEs in ARV-naïve patients enrolled in randomized trials or large observational cohorts appear to be declining with increased availability of better-tolerated and less toxic ART regimens and are generally less than 10%.¹⁻¹¹ In general, the overall benefits of ART outweigh its risks, and the risk of some abnormal laboratory findings (e.g., anemia, renal impairment) may be lower with ART than in its absence during HIV infection.

ARV drug-related AEs can vary in severity from mild to severe and life-threatening. Drug-related toxicity can be acute (occurring soon after a drug has been administered), subacute (occurring within 1 to 2 days of administration), or late (occurring after prolonged drug administration). For some ARV medications, pharmacogenetic markers associated with risk of early toxicity have been identified, but the only such screen in routine clinical use is HLA B*5701 as a marker for abacavir hypersensitivity.¹²⁻¹⁴ For selected children aged <3 years who require treatment with efavirenz, an additional pharmacogenetic marker, CYP2B6 genotype, should be assessed in an attempt to prevent toxicity (see [Efavirenz](#) in [Appendix A: Pediatric Antiretroviral Drug Information](#)).¹³⁻¹⁶ For a few other ARV drugs, known therapeutic ranges for plasma concentrations as determined by therapeutic drug monitoring (TDM) may indicate the need for dose reduction or modification of ART in patients experiencing AEs (see below and [Role of Therapeutic Drug Monitoring in Management of Pediatric HIV Infection](#)).

The most common acute and chronic AEs associated with ARV drugs or drug classes are presented in the [Management of Medication Toxicity or Intolerance](#) tables. The tables include information on common causative drugs, estimated frequency of occurrence, timing of symptoms, risk factors, potential preventive measures, and suggested clinical management strategies and provide selected references regarding these toxicities in pediatric patients.

Management

Management of medication-related toxicity should take into account its severity, the relative need for viral suppression, and the available ARV options. In general, mild and moderate toxicities do not require discontinuation of therapy or drug substitution. However, even mild AEs may have a negative impact on medication adherence and should be discussed before therapy is initiated, at regular provider visits, and at onset of any AEs. Common, self-limited AEs should be anticipated, and reassurance provided that many AEs will resolve after the first few weeks of ART. For example, when initiating therapy with boosted protease inhibitors (PIs), many patients experience gastrointestinal AEs such as nausea, vomiting, diarrhea, and abdominal pain. Instructing patients to take PIs with food may help minimize these AEs. Some patients may require antiemetics and antidiarrheal agents for symptom management. Central nervous system AEs are commonly encountered when initiating therapy with efavirenz. Symptoms can include dizziness, drowsiness, vivid dreams, or insomnia. Patients should be instructed to take efavirenz-containing regimens at bedtime, on an empty stomach, to help minimize these AEs. They should be advised that these AEs usually diminish in general within 2 to 4 weeks of initiating therapy in most people, but may persist for months in some, and may require a medication change.¹⁷⁻¹⁹ In addition, mild rash can be ameliorated with drugs such as antihistamines. For some moderate toxicities, using a drug in the same class as the one causing toxicity but with a different toxicity profile may be sufficient and discontinuation of all therapy may not be required.

In patients who experience unacceptable AEs from ART, every attempt should be made to identify the offending agent and to replace the drug with another effective agent as soon as possible.^{20,21} Many experts will stagger a planned interruption of a non-nucleoside reverse transcriptase inhibitor (NNRTI)-based regimen, stopping the NNRTI first and the dual nucleoside analogue reverse transcriptase backbone 7 to 14 days later because of the long half-life of NNRTI drugs. For patients who have a severe or life-threatening toxicity (e.g., hypersensitivity reaction—see [Hypersensitivity Reaction, Table 12I](#)), however, all components of the drug regimen should be stopped simultaneously, regardless of drug half-life. Once the offending drug or alternative cause for the AE has been determined, planning can begin for resumption of therapy with a new ARV regimen that does not contain the offending drug or with the original regimen, if the event is attributable to another cause. All drugs in the ARV regimen should then be started simultaneously, rather than one at a time with observation for AEs.

When therapy is changed because of toxicity or intolerance in a patient with virologic suppression, agents with different toxicity and side-effect profiles should be chosen, when possible.²²⁻²⁶ Clinicians should have comprehensive knowledge of the toxicity profile of each agent before selecting a new regimen. In the event of drug intolerance, changing a single drug in a multidrug regimen is permissible for patients whose viral loads are undetectable. However, substitution of a single active agent for a single drug in a failing multidrug regimen (e.g., a patient with virologic failure) is generally not recommended because of concern for development of resistance (see [Recognizing and Managing Antiretroviral Treatment Failure](#) in [Management of Children Receiving Antiretroviral Therapy](#)).

TDM may be used in the management of a child with mild or moderate toxicity if the toxicity is thought to be the result of a drug concentration exceeding the normal therapeutic range^{27,28} (see [Role of Therapeutic Drug Monitoring](#)). This is the only setting in which dose reduction would be considered appropriate management of drug toxicity, and even then, it should be used with caution; an expert in the management of pediatric HIV infection should be consulted.

To summarize, management strategies for drug intolerance include:

- Symptomatic treatment of mild-to-moderate transient side effects.
- If necessary, change from one drug to another drug to which a patient's virus is **susceptible** (such as changing to abacavir for zidovudine-related anemia or to **a PI or integrase strand transfer inhibitor** (INSTI) for efavirenz-related central nervous system symptoms).
- Change drug class, if necessary (e.g., from a PI to **an INSTI or a NNRTI** or vice versa) and if a patient's virus is **susceptible** to a drug in that class.
- Dose reduction only when drug **concentrations** are determined **to be above the therapeutic range**.

References

1. Sauvageot D, Schaefer M, Olson D, Pujades-Rodriguez M, O'Brien DP. Antiretroviral therapy outcomes in resource-limited settings for HIV-infected children <5 years of age. *Pediatrics*. 2010;125(5):e1039-1047. Available at <http://www.ncbi.nlm.nih.gov/pubmed/20385636>.
2. Buck WC, Kabue MM, Kazembe PN, Kline MW. Discontinuation of standard first-line antiretroviral therapy in a cohort of 1434 Malawian children. *J Int AIDS Soc*. 2010;13:31. Available at <http://www.ncbi.nlm.nih.gov/pubmed/20691049>.
3. Tukei VJ, Asiimwe A, Maganda A, et al. Safety and tolerability of antiretroviral therapy among HIV-infected children and adolescents in Uganda. *J Acquir Immune Defic Syndr*. 2012;59(3):274-280. Available at <http://www.ncbi.nlm.nih.gov/pubmed/22126740>.
4. Palmer M, Chersich M, Moultrie H, Kuhn L, Fairlie L, Meyers T. Frequency of stavudine substitution due to toxicity in children receiving antiretroviral treatment in sub-Saharan Africa. *AIDS*. 2013;27(5):781-785. Available at <http://www.ncbi.nlm.nih.gov/pubmed/23169331>.
5. Arpadi S, Shiau S, Strehlau R, et al. Metabolic abnormalities and body composition of HIV-infected children on Lopinavir or Nevirapine-based antiretroviral therapy. *Arch Dis Child*. 2013;98(4):258-264. Available at <http://www.ncbi.nlm.nih.gov/pubmed/23220209>.
6. Prendergast AJ. Complications of long-term antiretroviral therapy in HIV-infected children. *Arch Dis Child*. 2013;98(4):245-246. Available at <http://www.ncbi.nlm.nih.gov/pubmed/23413313>.
7. Purswani M, Patel K, Kopp JB, et al. Tenofovir treatment duration predicts proteinuria in a multiethnic United States Cohort of children and adolescents with perinatal HIV-1 infection. *Pediatr Infect Dis J*. 2013;32(5):495-500. Available at <http://www.ncbi.nlm.nih.gov/pubmed/23249917>.
8. Shubber Z, Calmy A, Andrieux-Meyer I, et al. Adverse events associated with nevirapine and efavirenz-based first-line antiretroviral therapy: a systematic review and meta-analysis. *AIDS*. 2013;27(9):1403-1412. Available at <http://www.ncbi.nlm.nih.gov/pubmed/23343913>.
9. Arrow Trial team, Kekitiinwa A, Cook A, et al. Routine versus clinically driven laboratory monitoring and first-line antiretroviral therapy strategies in African children with HIV (ARROW): a 5-year open-label randomised factorial trial. *Lancet*. 2013;381(9875):1391-1403. Available at <http://www.ncbi.nlm.nih.gov/pubmed/23473847>.
10. Barlow-Mosha L, Eckard AR, McComsey GA, Musoke PM. Metabolic complications and treatment of perinatally HIV-infected children and adolescents. *J Int AIDS Soc*. 2013;16:18600. Available at <http://www.ncbi.nlm.nih.gov/pubmed/23782481>.
11. Cohen S, Smit C, van Rossum AM, et al. Long-term response to combination antiretroviral therapy in HIV-infected children in the Netherlands registered from 1996 to 2012. *AIDS*. 2013;27(16):2567-2575. Available at <http://www.ncbi.nlm.nih.gov/pubmed/23842124>.
12. Lubomirov R, Colombo S, di Iulio J, et al. Association of pharmacogenetic markers with premature discontinuation of first-line anti-HIV therapy: an observational cohort study. *J Infect Dis*. 2011;203(2):246-257. Available at <http://www.ncbi.nlm.nih.gov/pubmed/21288825>.
13. Aceti A, Gianserra L, Lambiase L, Pennica A, Teti E. Pharmacogenetics as a tool to tailor antiretroviral therapy: A review. *World J Virol*. 2015;4(3):198-208. Available at <http://www.ncbi.nlm.nih.gov/pubmed/26279982>.
14. Asensi V, Collazos J, Valle-Garay E. Can antiretroviral therapy be tailored to each human immunodeficiency virus-

- infected individual? Role of pharmacogenomics. *World J Virol.* 2015;4(3):169-177. Available at <http://www.ncbi.nlm.nih.gov/pubmed/26279978>.
15. Sinxadi PZ, Leger PD, McIlleron HM, et al. Pharmacogenetics of plasma efavirenz exposure in HIV-infected adults and children in South Africa. *Br J Clin Pharmacol.* 2015;80(1):146-156. Available at <http://www.ncbi.nlm.nih.gov/pubmed/25611810>.
16. Haas DW, Ribaud HJ, Kim RB, et al. Pharmacogenetics of efavirenz and central nervous system side effects: an Adult AIDS Clinical Trials Group study. *AIDS.* 2004;18(18):2391-2400. Available at <http://www.ncbi.nlm.nih.gov/pubmed/15622315>.
17. Gazzard B, Duvivier C, Zagler C, et al. Phase 2 double-blind, randomized trial of etravirine versus efavirenz in treatment-naïve patients: 48-week results. *AIDS.* 2011;25(18):2249-2258. Available at <http://www.ncbi.nlm.nih.gov/pubmed/21881478>.
18. Nelson M, Stellbrink HJ, Podzamczar D, et al. A comparison of neuropsychiatric adverse events during 12 weeks of treatment with etravirine and efavirenz in a treatment-naïve, HIV-1-infected population. *AIDS.* 2011;25(3):335-340. Available at <http://www.ncbi.nlm.nih.gov/pubmed/21150563>.
19. Leutscher PD, Stecher C, Storgaard M, Larsen CS. Discontinuation of efavirenz therapy in HIV patients due to neuropsychiatric adverse effects. *Scand J of Infect Dis.* 2013;45(8):645-651. Available at <http://www.ncbi.nlm.nih.gov/pubmed/23427878>.
20. Elzi L, Marzolini C, Furrer H, et al. Treatment modification in human immunodeficiency virus-infected individuals starting combination antiretroviral therapy between 2005 and 2008. *Arch Intern Med.* 2010;170(1):57-65. Available at <http://www.ncbi.nlm.nih.gov/pubmed/20065200>.
21. Davidson I, Beardsell H, Smith B, et al. The frequency and reasons for antiretroviral switching with specific antiretroviral associations: the SWITCH study. *Antiviral Res.* 2010;86(2):227-229. Available at <http://www.ncbi.nlm.nih.gov/pubmed/20211651>.
22. Martinez E, Larrousse M, Llibre JM, et al. Substitution of raltegravir for ritonavir-boosted protease inhibitors in HIV-infected patients: the SPIRAL study. *AIDS.* 2010;24(11):1697-1707. Available at <http://www.ncbi.nlm.nih.gov/pubmed/20467288>.
23. McComsey G, Bhumbra N, Ma JF, Rathore M, Alvarez A, First Pediatric Switch S. Impact of protease inhibitor substitution with efavirenz in HIV-infected children: results of the First Pediatric Switch Study. *Pediatrics.* 2003;111(3):e275-281. Available at <http://www.ncbi.nlm.nih.gov/pubmed/12612284>.
24. Viergever RF, ten Berg MJ, van Solinge WW, Hoepelman AI, Gisolf EH. Changes in hematological parameters after switching treatment of HIV-infected patients from zidovudine to abacavir or tenofovir DF. *HIV Clin Trials.* 2009;10(2):125-128. Available at <http://www.ncbi.nlm.nih.gov/pubmed/19487183>.
25. Valantin MA, Bittar R, de Truchis P, et al. Switching the nucleoside reverse transcriptase inhibitor backbone to tenofovir disoproxil fumarate + emtricitabine promptly improves triglycerides and low-density lipoprotein cholesterol in dyslipidaemic patients. *J Antimicrob Chemother.* 2010;65(3):556-561. Available at <http://www.ncbi.nlm.nih.gov/pubmed/20053692>.
26. Mallolas J, Podzamczar D, Milinkovic A, et al. Efficacy and safety of switching from boosted lopinavir to boosted atazanavir in patients with virological suppression receiving a LPV/r-containing HAART: the ATAZIP study. *J Acquir Immune Defic Syndr.* 2009;51(1):29-36. Available at <http://www.ncbi.nlm.nih.gov/pubmed/19390327>.
27. van Luin M, Gras L, Richter C, et al. Efavirenz dose reduction is safe in patients with high plasma concentrations and may prevent efavirenz discontinuations. *J Acquir Immune Defic Syndr.* 2009;52(2):240-245. Available at <http://www.ncbi.nlm.nih.gov/pubmed/19593159>.
28. Pretorius E, Klinker H, Rosenkranz B. The role of therapeutic drug monitoring in the management of patients with human immunodeficiency virus infection. *Ther Drug Monit.* 2011;33(3):265-274. Available at <http://www.ncbi.nlm.nih.gov/pubmed/21566505>.

Table 12a. Antiretroviral-Therapy-Associated Adverse Effects and Management Recommendations—Central Nervous System Toxicity (Last updated March 1, 2016; last reviewed March 1, 2016) (page 1 of 3)

Adverse Effects	Associated ARVs	Onset/Clinical Manifestations	Estimated Frequency	Risk Factors	Prevention/Monitoring	Management
Global CNS Depression	LPV/r oral solution (contains both ethanol and propylene glycol as excipients)	<p><u>Onset:</u></p> <ul style="list-style-type: none"> • 1–6 days after starting LPV/r <p><u>Presentation</u></p> <p><i>Neonates/Preterm Infants:</i></p> <ul style="list-style-type: none"> • Global CNS depression (e.g., abnormal EEG, altered state of consciousness, somnolence) • Non-CNS-associated toxicity may include cardiac toxicity and respiratory complications. 	<p>Exact frequency of ethanol and propylene glycol-associated toxicity unknown in neonates receiving LPV/r oral solution.</p>	<p>Prematurity</p> <p>Low birth weight</p> <p>Age <14 days (whether premature or term)</p>	<p>Avoid use of LPV/r until a postmenstrual age of 42 weeks and a postnatal age ≥14 days.</p>	<p>Discontinue LPV/r; symptoms should resolve in 1–5 days.</p> <p>If needed, reintroduction of LPV/r can be considered once outside the vulnerable period (i.e., postmenstrual age of 42 weeks and a postnatal age ≥14 days).</p>
Neuropsychiatric Symptoms and Other CNS Manifestations	EFV	<p><u>Onset:</u></p> <ul style="list-style-type: none"> • 1–2 days after initiating treatment • Many symptoms subside or diminish by 2–4 weeks, but may persist in a significant proportion of patients. In one report, 37% experienced persistent symptoms at 12 months and in another, half of discontinuations occurred after 12 months. <p><u>Presentation (May Include One or More of the Following)</u></p> <p>Neuropsychiatric Symptoms:</p> <ul style="list-style-type: none"> • Abnormal dreams • Psychosis • Suicidal ideation or attempted/completed suicide • Seizures (including absence seizures) or decreased seizure threshold 	<p>Variable, depending on age, symptom, assessment method</p> <p><u>Children:</u></p> <ul style="list-style-type: none"> • 24% for any EFV-related CNS manifestations in 1 case series with 18% requiring drug discontinuation • 9% incidence of new-onset seizures reported in 1 study in children aged <36 months, in two of the children the seizures had alternative causes. <p><u>Adults:</u></p> <ul style="list-style-type: none"> • 30% incidence for any CNS manifestations of any severity. • 6% incidence for EFV-related severe CNS manifestations including suicidality. 	<p>Insomnia associated with elevated EFV trough concentration ≥4 mcg/mL</p> <p>Presence of CYP450 polymorphisms that decrease EFV metabolism (CYP2B6 516 TT genotype)</p> <p>Prior history of psychiatric illness or use of psychoactive drugs</p>	<p>Administer EFV on an empty stomach, preferably at bedtime.</p> <p>Use with caution in the presence of psychiatric illness including depression or suicidal thoughts or with concomitant use of psychoactive drugs.</p> <p>TDM can be considered in the context of a child with mild or moderate toxicity possibly attributable to a particular ARV agent (see Role of Therapeutic Drug Monitoring in Management of Treatment Failure).</p>	<p>Obtain EFV trough concentration if symptoms excessive or persistent. If EFV trough concentration >4 mcg/mL, strongly consider drug substitution if suitable alternative exists. Alternatively, consider dose reduction with repeat TDM and dose adjustment (with expert pharmacologist input).</p> <p>In a small study, cyproheptadine was shown to reduce short-term incidence of neuropsychiatric effects in adults receiving EFV, but data are lacking in children and no recommendation can be made for its use at this time.</p>

Table 12a. Antiretroviral-Therapy-Associated Adverse Effects and Management Recommendations—Central Nervous System (CNS) Toxicity (Last updated March 1, 2016; last reviewed March 1, 2016) (page 2 of 3)

Adverse Effects	Associated ARVs	Onset/Clinical Manifestations	Estimated Frequency	Risk Factors	Prevention/Monitoring	Management
Neuropsychiatric Symptoms and Other CNS Manifestations, continued	EFV, continued	<p><i>Other CNS Manifestations:</i></p> <ul style="list-style-type: none"> • Dizziness • Somnolence • Insomnia or poor sleep quality • Impaired concentration <p>Note: Some CNS side effects (e.g., impaired concentration, abnormal dreams, or sleep disturbances) may be more difficult to assess in children.</p>	However, evidence is conflicting about whether EFV use increases the incidence of suicidality.			
	RPV	<p><u>Presentation</u></p> <p><i>Neuropsychiatric Symptoms:</i></p> <ul style="list-style-type: none"> • Depressive disorders • Suicidal ideation • Abnormal dreams/nightmare <p><i>Other CNS Manifestations:</i></p> <ul style="list-style-type: none"> • Headache • Dizziness • Insomnia 	<p><u>In Adults:</u></p> <ul style="list-style-type: none"> • CNS/neuropsychiatric adverse events of all severity grades were reported in 43% of patients at 96 weeks (mostly Grade 1). Depressive disorders of all severity grades were reported in 9% of patients, and were severe requiring RPV discontinuation in 1% of patients. <p><u>In Children:</u></p> <ul style="list-style-type: none"> • Depressive disorders of all severity grades were reported in 19.4% of pediatric patients aged 12 years to 17 years. Severe depressive disorders were reported in 5.6% of patients, including a suicide attempt in 1 subject. 	Prior history of neuropsychiatric illness	Monitor carefully for depressive disorders and other CNS symptoms.	Consider drug substitution in case of severe symptoms.
	RAL	<p><u>Presentation:</u></p> <ul style="list-style-type: none"> • Increased psychomotor activity • Headaches • Insomnia • Depression 	<p><u>Children:</u></p> <ul style="list-style-type: none"> • Increased psychomotor activity reported in one child. <p><u>Adults:</u></p> <ul style="list-style-type: none"> • Headache • Insomnia (<5% in adult trials) 	<p>Elevated RAL concentrations</p> <p>Co-treatment with TDF or PPI</p> <p>Prior history of insomnia or depression</p>	<p>Prescreen for psychiatric symptoms.</p> <p>Monitor carefully for CNS symptoms.</p> <p>Use with caution in the presence of drugs that increase RAL concentration.</p>	Consider drug substitution (RAL or co-administered drug) in case of severe insomnia or other neuropsychiatric symptoms.

Table 12a. Antiretroviral-Therapy-Associated Adverse Effects and Management Recommendations—Central Nervous System (CNS) Toxicity (Last updated March 1, 2016; last reviewed March 1, 2016) (page 3 of 3)

Adverse Effects	Associated ARVs	Onset/Clinical Manifestations	Estimated Frequency	Risk Factors	Prevention/Monitoring	Management
	DTG	<p><u>Onset:</u></p> <ul style="list-style-type: none"> • 7–30 days after initiating drug <p><u>Presentation</u></p> <p><i>Neuropsychiatric Symptoms:</i></p> <ul style="list-style-type: none"> • Depression or exacerbation of preexisting depression • Anxiety • Suicidal ideation attempt, behavior, or completion <p><i>Other CNS Manifestations (Generally Mild):</i></p> <ul style="list-style-type: none"> • Insomnia • Dizziness • Headache 	<p><u>Adults</u></p> <ul style="list-style-type: none"> • Exact frequency of neuropsychiatric symptoms is unknown; case reports of 4 adult patients. Headache, insomnia, and dizziness are common, reported in up to 10% of patients. Less than 1% patients experienced more severe symptoms. 	Pre-existing depression or other psychiatric illness	Use with caution in the presence of psychiatric illness especially depression	<p>For severe neuropsychiatric symptoms, consider discontinuation of DTG if suitable alternative exists.</p> <p>Discontinuation resulted in resolution of neuropsychiatric symptom in 3 out of 4 patients (in the 4th patient, symptoms resolved slowly despite DTG continuation).</p> <p>For mild symptoms, continue DTG and counsel patient that symptoms will likely resolve with time.</p>
Intracranial Hemorrhage	TPV	<p><u>Onset:</u></p> <ul style="list-style-type: none"> • 7–513 days after starting TPV 	<p><u>Children:</u></p> <ul style="list-style-type: none"> • No cases of ICH reported in children. <p><u>Adults:</u></p> <ul style="list-style-type: none"> • In premarket approval data in adults, 0.23/100 py or 0.04–0.22/100 py in a retrospective review of 2 large patient databases. 	Unknown; prior history of bleeding disorder or risk factors for bleeding present in most patients in case series reported.	Administer TPV with caution in patients with bleeding disorder, known intracranial lesions, or recent neurosurgery.	Discontinue TPV if ICH is suspected or confirmed.
Cerebellar Ataxia	RAL	<p><u>Onset:</u></p> <ul style="list-style-type: none"> • As early as 3 days after starting RAL <p><u>Presentation:</u></p> <ul style="list-style-type: none"> • Tremor • Dysmetria • Ataxia 	Two cases reported in adults during post-marketing period.	Unknown; a speculated mechanism may include recent treatment with ATV with residual UGT1A1 enzyme inhibition and increased RAL serum concentration.	Use with caution with ATV or other drugs that cause strong inhibition of UGT1A1 enzyme.	Consider drug discontinuation. RAL reintroduction can be considered if predisposing factor (e.g., drug-drug interaction) identified and removed.

Key to Acronyms: ARV = antiretroviral; ATV = atazanavir; CNS = central nervous system; CYP = cytochrome P; **DTG = dolutegravir**; EEG = electroencephalogram; EFV = efavirenz; ICH = intracranial hemorrhage; LPV/r = ritonavir-boosted lopinavir; PPI = proton pump inhibitor; py = patient years; RAL = raltegravir; RPV = rilpivirine; TDF = tenofovir disoproxil fumarate; TDM = therapeutic drug monitoring; TPV = tipranavir; UGT = uridine diphosphate-glucurononyl transferase

References

1. FDA. FDA Drug Safety Communication: Serious health problems seen in premature babies given Kaletra (lopinavir/ritonavir) oral solution. Accessed 4/28/14. 2011. Available at <http://www.fda.gov/Drugs/DrugSafety/ucm246002.htm>.
2. Gutierrez F, Navarro A, Padilla S, et al. Prediction of neuropsychiatric adverse events associated with long-term efavirenz therapy, using plasma drug level monitoring. *Clin Infect Dis*. 2005;41(11):1648-1653. Available at <http://www.ncbi.nlm.nih.gov/pubmed/16267739>.
3. Haas DW, Ribaud HJ, Kim RB, et al. Pharmacogenetics of efavirenz and central nervous system side effects: an Adult AIDS Clinical Trials Group study. *AIDS*. 2004;18(18):2391-2400. Available at <http://www.ncbi.nlm.nih.gov/pubmed/15622315>.
4. Waters L, Fisher M, Winston A, et al. A phase IV, double-blind, multicentre, randomized, placebo-controlled, pilot study to assess the feasibility of switching individuals receiving efavirenz with continuing central nervous system adverse events to etravirine. *AIDS*. 2011;25(1):65-71. Available at <http://www.ncbi.nlm.nih.gov/pubmed/21099666>.
5. Puthanakit T, Tanpaiboon P, Aupibul L, Cressey TR, Sirisanthana V. Plasma efavirenz concentrations and the association with CYP2B6-516G >T polymorphism in HIV-infected Thai children. *Antivir Ther*. 2009;14(3):315-320. Available at <http://www.ncbi.nlm.nih.gov/pubmed/19474465>.
6. Cabrera Figueroa S, Fernandez de Gatta M, Hernandez Garcia L, et al. The convergence of therapeutic drug monitoring and pharmacogenetic testing to optimize efavirenz therapy. *Ther Drug Monit*. 2010;32(5):579-585. Available at <http://www.ncbi.nlm.nih.gov/pubmed/20720517>.
7. Tepller H, Brown DD, Leavitt RY, et al. Long-term safety from the raltegravir clinical development program. *Current HIV research*. 2011;9(1):40-53. Available at <http://www.ncbi.nlm.nih.gov/pubmed/21198432>.
8. Reiss KA, Bailey JR, Pham PA, Gallant JE. Raltegravir-induced cerebellar ataxia. *AIDS*. 2010;24(17):2757. Available at <http://www.ncbi.nlm.nih.gov/pubmed/20980871>.
9. Strehlau R, Martens L, Coovadia A, et al. Absence seizures associated with efavirenz initiation. *Pediatr Infect Dis J*. 2011;30(11):1001-1003. Available at <http://www.ncbi.nlm.nih.gov/pubmed/21633320>.
10. Rakhmanina NY, van den Anker JN, Soldin SJ, van Schaik RH, Mordwinkin N, Neely MN. Can therapeutic drug monitoring improve pharmacotherapy of HIV infection in adolescents? *Ther Drug Monit*. 2010;32(3):273-281. Available at <http://www.ncbi.nlm.nih.gov/pubmed/20445485>.
11. Cattaneo D, Ripamonti D, Baldelli S, Cozzi V, Conti F, Clementi E. Exposure-related effects of atazanavir on the pharmacokinetics of raltegravir in HIV-1-infected patients. *Ther Drug Monit*. 2010;32(6):782-786. Available at <http://www.ncbi.nlm.nih.gov/pubmed/20926993>.
12. Chan-Tack KM, Struble KA, Birnkrant DB. Intracranial hemorrhage and liver-associated deaths associated with tipranavir/ritonavir: review of cases from the FDA's Adverse Event Reporting System. *AIDS Patient Care STDS*. 2008;22(11):843-850. Available at <http://www.ncbi.nlm.nih.gov/pubmed/19025478>.
13. Shubber Z, Calmy A, Andrieux-Meyer I, et al. Adverse events associated with nevirapine and efavirenz-based first-line antiretroviral therapy: a systematic review and meta-analysis. *AIDS*. 2013;27(9):1403-1412. Available at <http://www.ncbi.nlm.nih.gov/pubmed/23343913>.
14. van Dijk JH, Sutcliffe CG, Hamangaba F, Bositis C, Watson DC, Moss WJ. Effectiveness of efavirenz-based regimens in young HIV-infected children treated for tuberculosis: a treatment option for resource-limited settings. *PLoS One*. 2013;8(1):e55111. Available at <http://www.ncbi.nlm.nih.gov/pubmed/23372824>.
15. Cohen CJ, Molina JM, Cassetti I, et al. Week 96 efficacy and safety of rilpivirine in treatment-naïve, HIV-1 patients in two Phase III randomized trials. *AIDS*. 2013;27(6):939-950. Available at <http://www.ncbi.nlm.nih.gov/pubmed/23211772>.
16. Nachman S, al. E. IMPAACT P1066: raltegravir (RAL) safety and efficacy in HIV infected (+) youth two to 18 years of age through week 48. Abstract no. TUAB0205. 19th International AIDS Conference; 2012; Washington, DC.
17. Madeddu G, Menzaghi B, Ricci E, et al. Raltegravir central nervous system tolerability in clinical practice: results from a multicenter observational study. *AIDS*.

2012;26(18):2412-2415. Available at <http://www.ncbi.nlm.nih.gov/pubmed/23032413>.

18. Dabaghzadeh F, Ghaeli P, Khalili H, et al. Cyproheptadine for prevention of neuropsychiatric adverse effects of efavirenz: a randomized clinical trial. *AIDS Patient Care STDS*. 2013;27(3):146-154. Available at <http://www.ncbi.nlm.nih.gov/pubmed/23442031>.
19. Mollan KR, Smurzynski M, Eron JJ, et al. Association between efavirenz as initial therapy for HIV-1 infection and increased risk for suicidal ideation or attempted or completed suicide: an analysis of trial data. *Ann Intern Med*. 2014;161(1):1-10. Available at <http://www.ncbi.nlm.nih.gov/pubmed/24979445>.
20. Mills AM, Antinori A, Clotet B, et al. Neurological and psychiatric tolerability of rilpivirine (TMC278) vs. efavirenz in treatment-naïve, HIV-1-infected patients at 48 weeks. *HIV Med*. 2013;14(7):391-400. Available at <http://www.ncbi.nlm.nih.gov/pubmed/23298380>.
21. Napoli AA, Wood JJ, Coumbis JJ, Soitkar AM, Seekins DW, Tilson HH. No evident association between efavirenz use and suicidality was identified from a disproportionality analysis using the FAERS database. *J Int AIDS Soc*. 2014;17:19214. Available at <http://www.ncbi.nlm.nih.gov/pubmed/25192857>.
22. Nkhoma Eea. Using Real World Data to Assess the Risk of Suicidality among Patients Initiating an Efavirenz-containing Regimen versus an Efavirenz-free Antiretroviral Regimen. Infectious Disease Society of America: ID Week 2014; 2014; Philadelphia, PA.
23. Leutscher PD, Stecher C, Storgaard M, Larsen CS. Discontinuation of efavirenz therapy in HIV patients due to neuropsychiatric adverse effects. *Scandinavian journal of infectious diseases*. 2013;45(8):645-651. Available at <http://www.ncbi.nlm.nih.gov/pubmed/23427878>.
24. Group ES, Puls R, Amin J, et al. Efficacy of 400 mg efavirenz versus standard 600 mg dose in HIV-infected, antiretroviral-naïve adults (ENCORE1): a randomised, double-blind, placebo-controlled, non-inferiority trial. *Lancet*. 2014;383(9927):1474-1482. Available at <http://www.ncbi.nlm.nih.gov/pubmed/24522178>.
25. Smith C, Ryom L, Monforte A, et al. Lack of association between use of efavirenz and death from suicide: evidence from the D:A:D study. *J Int AIDS Soc*. 2014;17(4 Suppl 3):19512. Available at <http://www.ncbi.nlm.nih.gov/pubmed/25394021>.
26. Ford N, Shubber Z, Pozniak A, et al. Comparative Safety and Neuropsychiatric Adverse Events Associated With Efavirenz Use in First-Line Antiretroviral Therapy: A Systematic Review and Meta-Analysis of Randomized Trials. *J Acquir Immune Defic Syndr*. 2015;69(4):422-429. Available at <http://www.ncbi.nlm.nih.gov/pubmed/25850607>.
27. Mills A, Garner W, Pozniak A, et al. Patient-Reported Symptoms Over 48 Weeks in a Randomized, Open-Label, Phase IIb Non-Inferiority Trial of Adults with HIV Switching to Co-Formulated Elvitegravir, Cobicistat, Emtricitabine, and Tenofovir DF versus Continuation of Non-Nucleoside Reverse Transcriptase Inhibitor with Emtricitabine and Tenofovir DF. *Patient*. 2015;8(4):359-371. Available at <http://www.ncbi.nlm.nih.gov/pubmed/26045359>.
28. García-Navarro C, Jiménez de Ory S, Navarro Gómez ML, et al. Sleep disturbances in a cohort of hiv-infected children and adolescents on antiretroviral treatment. 22nd Conference on Retroviruses and Opportunistic Infections 2015; Seattle, WA.
29. Kheloufi F, Allemand J, Mokhtari S, Default A. Psychiatric disorders after starting dolutegravir: report of four cases. *AIDS*. 2015;29(13):1723-1725.
30. Food and Drug Administration. Edurant Package Insert. Available at http://www.accessdata.fda.gov/drugsatfda_docs/label/2015/202022s0081bledt.pdf 2015.
31. Walmsley S, Baumgarten A, Berenguer J, et al. Dolutegravir Plus Abacavir/Lamivudine for the Treatment of HIV-1 Infection in Antiretroviral Therapy-Naïve Patients: Week 96 and Week 144 Results from the SINGLE Randomized Clinical Trial. *J Acquir Immune Defic Syndr*. 2015. Available at <http://www.ncbi.nlm.nih.gov/pubmed/26262777>.

Table 12b. Antiretroviral-Therapy-Associated Adverse Effects and Management Recommendations—Dyslipidemia

(Last updated March 1, 2016; last reviewed March 1, 2016) (page 1 of 2)

Adverse Effects	Associated ARVs	Onset/Clinical Manifestations	Estimated Frequency	Risk Factors	Prevention/Monitoring	Management
Dyslipidemia	<p><u>PIs:</u></p> <ul style="list-style-type: none"> All PIs, especially RTV-boosted PIs; lower incidence reported with DRV/r and ATV with or without RTV. <p><u>NRTIs:</u></p> <ul style="list-style-type: none"> Especially d4T <p><u>NNRTIs:</u></p> <ul style="list-style-type: none"> EFV > NVP, RPV, and ETR 	<p><u>Onset:</u></p> <ul style="list-style-type: none"> As early as 2 weeks to months after beginning therapy <p><u>Presentation</u></p> <p><u>PIs:</u></p> <ul style="list-style-type: none"> ↑LDL-C, TC, and TG <p><u>NNRTIs:</u></p> <ul style="list-style-type: none"> ↑LDL-C, TC, and HDL-C <p><u>NRTIs:</u></p> <ul style="list-style-type: none"> ↑LDL-C, TC, and TG 	<p>Reported frequency varies with specific ARV regimen, duration of ART and specific laboratory parameters used to diagnose lipid abnormalities.</p> <p>10% to 20% in young children receiving LPV/RTV</p> <p>40% to 75% of older children and adolescents with prolonged ART history will have lipid abnormalities.</p> <p>In studies of treatment naive adults, 38% and 32% receiving EVG/COBI/FTC/TAF developed abnormal fasting TC and LDL-C (respectively) after 48 weeks compared with 21% and 20% receiving EVG/COBI/FTC/TDF; difference mainly attributable to TAF</p> <p>In 48 adolescents treated with EVG/COBI/FTC/TAF median change from baseline to</p>	<p>Advanced-stage HIV disease</p> <p>High-fat, high-cholesterol diet</p> <p>Lack of exercise</p> <p>Obesity</p> <p>Hypertension</p> <p>Smoking</p> <p>Family history of dyslipidemia or premature CVD</p> <p>Metabolic syndrome</p> <p>Fat maldistribution</p>	<p><u>Prevention:</u></p> <ul style="list-style-type: none"> Low-fat diet Exercise Smoking-prevention counseling <p><u>Monitoring^a</u></p> <p><i>Adolescents and Adults:</i></p> <ul style="list-style-type: none"> Monitor 12-hour FLP, which includes TC, HDL-C, non-HDL-C, LDL-C, and TG, every 6–12 months. Obtain FLPs twice (>2 weeks but ≤3 months apart, average results) before initiating or changing lipid-lowering therapy. <p><i>Children (Aged ≥2 Years) without Lipid Abnormalities or Additional Risk Factors:</i></p> <ul style="list-style-type: none"> Obtain non-fasting screening lipid profiles before initiating or changing therapy and then, if levels are stable, every 6–12 months. If TG or LDL-C is elevated, obtain fasting blood tests. <p><i>Children with Lipid Abnormalities and/or Additional Risk Factors:</i></p> <ul style="list-style-type: none"> Obtain 12-hour FLP before initiating or changing therapy and every 6 months thereafter (more often if indicated). <p><i>Children Receiving Lipid-Lowering Therapy with Statins or Fibrates:</i></p> <ul style="list-style-type: none"> Obtain 12-hour FLP, LFTs, and CK at 4 and 8 weeks, and 3 	<p>Assessment of additional CVD risk factors should be done in all patients. HIV-infected patients are considered to be at moderate risk of CVD.^b</p> <p>Counsel on lifestyle modification, dietary interventions (e.g., a diet low in saturated fat, cholesterol, and refined sugars particularly in case of ↑TG, elimination of trans fat, physical activity, smoking cessation) for an adequate trial period (3–6 months). Consider consultation with dietician.</p> <p>If receiving d4T, it should be discontinued. If receiving PI-based ART, consider switching to a new PI-sparing ART regimen or PI-based regimen containing boosted ATV or DRV, which are less likely to cause lipid abnormalities.</p> <p>Consider lipid-lowering therapy in consultation with a lipid specialist if ≥6-month trial of lifestyle modification fails.</p> <p>Some experts suggest treatment in children receiving ARV drugs at cut points recommended by NHLBI cardiovascular risk reduction guidelines for children aged ≥10 years: LDL-C ≥190 mg/dL, regardless of additional risk factors; LDL-C ≥160 mg/dL or LDL-C ≥130 mg/dL based on presence of additional risk factors and risk conditions.^b</p> <p>The minimal goal of therapy should be to achieve and maintain a LDL-C value below 130 mg/dL.</p>

Table 12b. Antiretroviral-Therapy-Associated Adverse Effects and Management Recommendations—Dyslipidemia

(Last updated March 1, 2016; last reviewed March 1, 2016) (page 2 of 2)

Adverse Effects	Associated ARVs	Onset/Clinical Manifestations	Estimated Frequency	Risk Factors	Prevention/Monitoring	Management
			weeks 24 and 36 were 26 mg/dl and 36 mg/dl, respectively for fasting TC, and 10 mg/dl and 17 mg/dl, respectively for direct LDL-C.		<p>months after starting lipid therapy.</p> <ul style="list-style-type: none"> If minimal alterations in AST, ALT, and CK, monitor every 3–4 months in the first year and every 6 months thereafter (or as clinically indicated). Repeat FLPs 4 weeks after increasing doses of antihyperlipidemic agents. 	<p><u>Initiate Drug Therapy Promptly in Patients with Fasting TG ≥ 500 mg/dL:</u></p> <p>Statins such as pravastatin, atorvastatin, or rosuvastatin.^c Ezetimibe can be considered in addition to statins.^d Statin-related toxicities include liver enzyme elevation and myopathy, and risk may be increased by drug interactions with ART, particularly PIs.^c Risks must be weighed against potential benefits.</p> <p>Fibrates (gemfibrozil and fenofibrate) and N-3 PUFAs derived from fish oils may be used as alternative agents for adults with \uparrow TG but are not approved for use in children. The long-term risks of lipid abnormalities in children receiving ART are unclear. However, persistent dyslipidemia in children may lead to premature CVD.</p>

^a Given the burden of collecting fasting blood samples, some practitioners routinely measure cholesterol and triglycerides from non-fasting blood samples and follow up abnormal values with a test done in the fasted state.

^b Refer to NHLBI guidelines at http://www.nhlbi.nih.gov/guidelines/cvd_ped/summary.htm#chap9.

^c The risks of new treatment-related toxicities and virologic failure that could occur with changes in therapy must be weighed against the potential risk of drug interactions and toxicities associated with the use of lipid-lowering agents.

^d Statins (HMG-CoA reductase inhibitors) are contraindicated in pregnancy (potentially teratogenic) and should not be used in patients who may become pregnant. Multiple drug interactions exist between ARV drugs and statins (exception pravastatin, which is not dependent on CYP3A4 for metabolism). Pravastatin, atorvastatin, rosuvastatin (Crestor®), fluvastatin, and ezetimibe (Zetia®) are approved for use in children aged ≥ 10 years. For additional information, see the PI, NNRTI, NRTI, and INSTI Drug Interactions Tables in the [Guidelines for the use of Antiretroviral Agents in HIV-1-Infected Adults and Adolescents](#).

Key to Acronyms: ALT = alanine aminotransferase; ART = antiretroviral therapy; ARV = antiretroviral; AST = aspartate aminotransferase; ATV = atazanavir; CK = creatine kinase; CVD = cardiovascular disease; CYP3A4 = cytochrome P450 3A4; d4T = stavudine; DRV = darunavir; DRV/r = ritonavir-boosted darunavir; EFV = efavirenz; ETR = etravirine; FLP = fasting lipid

profile; HDL-C = high-density lipoprotein cholesterol; LDL-C = low-density lipoprotein cholesterol; LFT = liver function test; LPV = lopinavir; NHLBI = National Heart, Lung, and Blood Institute; NNRTI = non-nucleoside reverse transcriptase inhibitor; NRTI = nucleoside reverse transcriptase inhibitor; NVP = nevirapine; PI = protease inhibitor; PUFA = polyunsaturated fatty acid; RPV = rilpivirine; RTV = ritonavir; TC = total cholesterol; TG = triglyceride. lopinavir; NHLBI = National Heart, Lung, and Blood Institute; NNRTI = non-nucleoside reverse transcriptase inhibitor; NRTI = nucleoside reverse transcriptase inhibitor; NVP = nevirapine; PI = protease inhibitor; PUFA = polyunsaturated fatty acid; RPV = rilpivirine; RTV = ritonavir; TC = total cholesterol; TG = triglyceride

References

1. Belay B, Belamarich PF, Tom-Revzon C. The use of statins in pediatrics: knowledge base, limitations, and future directions. *Pediatrics*. 2007;119(2):370-380. Available at <http://www.ncbi.nlm.nih.gov/pubmed/17272627>.
2. McComsey G, Bhumbra N, Ma JF, Rathore M, Alvarez A, First Pediatric Switch S. Impact of protease inhibitor substitution with efavirenz in HIV-infected children: results of the First Pediatric Switch Study. *Pediatrics*. 2003;111(3):e275-281. Available at <http://www.ncbi.nlm.nih.gov/pubmed/12612284>.
3. Engler MM, Engler MB, Malloy MJ, Paul SM, Kulkarni KR, Mietus-Snyder ML. Effect of docosahexaenoic acid on lipoprotein subclasses in hyperlipidemic children (the EARLY study). *Am J Cardiol*. 2005;95(7):869-871. Available at <http://www.ncbi.nlm.nih.gov/pubmed/15781019>.
4. Vigano A, Aldrovandi GM, Giacomet V, et al. Improvement in dyslipidaemia after switching stavudine to tenofovir and replacing protease inhibitors with efavirenz in HIV-infected children. *Antivir Ther*. 2005;10(8):917-924. Available at <http://www.ncbi.nlm.nih.gov/pubmed/16430197>.
5. Carter RJ, Wiener J, Abrams EJ, et al. Dyslipidemia among perinatally HIV-infected children enrolled in the PACTS-HOPE cohort, 1999-2004: a longitudinal analysis. *J Acquir Immune Defic Syndr*. 2006;41(4):453-460. Available at <http://www.ncbi.nlm.nih.gov/pubmed/16652053>.
6. Tassiopoulos K, Williams PL, Seage GR, 3rd, et al. Association of hypercholesterolemia incidence with antiretroviral treatment, including protease inhibitors, among perinatally HIV-infected children. *J Acquir Immune Defic Syndr*. 2008;47(5):607-614. Available at <http://www.ncbi.nlm.nih.gov/pubmed/18209684>.
7. Aldrovandi GM, Lindsey JC, Jacobson DL, et al. Morphologic and metabolic abnormalities in vertically HIV-infected children and youth. *AIDS*. 2009;23(6):661-672. Available at <http://www.ncbi.nlm.nih.gov/pubmed/19279441>.
8. Chantry CJ, Hughes MD, Alvero C, et al. Lipid and glucose alterations in HIV-infected children beginning or changing antiretroviral therapy. *Pediatrics*. 2008;122(1):e129-138. Available at <http://www.ncbi.nlm.nih.gov/pubmed/18519448>.
9. Rhoads MP, Lanigan J, Smith CJ, Lyall EG. Effect of specific ART drugs on lipid changes and the need for lipid management in children with HIV. *J Acquir Immune Defic Syndr*. 2011;57(5):404-412. Available at <http://www.ncbi.nlm.nih.gov/pubmed/21499114>.
10. Jacobson DL, Williams P, Tassiopoulos K, Melvin A, Hazra R, Farley J. Clinical management and follow-up of hypercholesterolemia among perinatally HIV-infected children enrolled in the PACTG 219C study. *J Acquir Immune Defic Syndr*. 2011;57(5):413-420. Available at <http://www.ncbi.nlm.nih.gov/pubmed/21602698>.
11. O'Gorman CS, O'Neill MB, Conwell LS. Considering statins for cholesterol-reduction in children if lifestyle and diet changes do not improve their health: a review of the risks and benefits. *Vasc Health Risk Manage*. 2011;7:1-14. Available at <http://www.ncbi.nlm.nih.gov/pubmed/21339908>.
12. Estrada V, Portilla J. Dyslipidemia related to antiretroviral therapy. *AIDS Reviews*. 2011;13(1):49-56. Available at <http://www.ncbi.nlm.nih.gov/pubmed/21412389>.
13. Feeney ER, Mallon PW. HIV and HAART-Associated Dyslipidemia. *Open Cardiovasc Med J*. 2011;5:49-63. Available at <http://www.ncbi.nlm.nih.gov/pubmed/21643501>.
14. Dube MP, Cadden JJ. Lipid metabolism in treated HIV Infection. *Best Pract Res Clin Endocrinol Metab*. 2011;25(3):429-442. Available at <http://www.ncbi.nlm.nih.gov/pubmed/21663837>.
15. Singh S, Willig JH, Mugavero MJ, et al. Comparative Effectiveness and Toxicity of Statins Among HIV-Infected Patients. *Clin Infect Dis*. 2011;52(3):387-395.

Available at <http://www.ncbi.nlm.nih.gov/pubmed/21189273>.

16. Expert Panel on Integrated Guidelines for Cardiovascular Health and Risk Reduction in Children and Adolescents. Integrated Guidelines for Cardiovascular Health and Risk Reduction in Children and Adolescents. The Report of the Expert Panel. 2011. Available at http://www.nhlbi.nih.gov/guidelines/cvd_ped/summary.htm.
17. Food and Drug Administration. FDA Drug Safety Communication: Interactions between certain HIV or hepatitis C drugs and cholesterol-lowering statin drugs can increase the risk of muscle injury. 2012. Available at <http://www.fda.gov/Drugs/DrugSafety/ucm293877.htm>.
18. Langat A, Benki-Nugent S, Wamalwa D, et al. Lipid changes in Kenyan HIV-1-infected infants initiating highly active antiretroviral therapy by 1 year of age. *Pediatr Infect Dis J*. 2013;32(7):e298-304. Available at <http://www.ncbi.nlm.nih.gov/pubmed/23385950>.
19. Casado JL, de Los Santos I, Del Palacio M, et al. Lipid-lowering effect and efficacy after switching to etravirine in HIV-infected patients with intolerance to suppressive HAART. *HIV Clin Trials*. 2013;14(1):1-9. Available at <http://www.ncbi.nlm.nih.gov/pubmed/23372109>.
20. Calza L, Manfredi R, Colangeli V, et al. Two-year treatment with rosuvastatin reduces carotid intima-media thickness in HIV type 1-infected patients receiving highly active antiretroviral therapy with asymptomatic atherosclerosis and moderate cardiovascular risk. *AIDS Res Hum Retroviruses*. 2013;29(3):547-556. Available at <http://www.ncbi.nlm.nih.gov/pubmed/23098891>.
21. Lazzaretti RK, Kuhmmer R, Sprinz E, Polanczyk CA, Ribeiro JP. Dietary intervention prevents dyslipidemia associated with highly active antiretroviral therapy in human immunodeficiency virus type 1-infected individuals: a randomized trial. *J Am Coll Cardiol*. 2012;59(11):979-988. Available at <http://www.ncbi.nlm.nih.gov/pubmed/22402068>.
22. Peters BS, Wierzbicki AS, Moyle G, Nair D, Brockmeyer N. The effect of a 12-week course of omega-3 polyunsaturated fatty acids on lipid parameters in hypertriglyceridemic adult HIV-infected patients undergoing HAART: a randomized, placebo-controlled pilot trial. *Clin Ther*. 2012;34(1):67-76. Available at <http://www.ncbi.nlm.nih.gov/pubmed/22212377>.
23. Oliveira JM, Rondo PH. Omega-3 fatty acids and hypertriglyceridemia in HIV-infected subjects on antiretroviral therapy: systematic review and meta-analysis. *HIV Clin Trials*. 2011;12(5):268-274. Available at <http://www.ncbi.nlm.nih.gov/pubmed/22180524>.
24. Strehlau R, Coovadia A, Abrams EJ, et al. Lipid profiles in young HIV-infected children initiating and changing antiretroviral therapy. *J Acquir Immune Defic Syndr*. 2012;60(4):369-376. Available at <http://www.ncbi.nlm.nih.gov/pubmed/22134152>.
25. Hazra R, Cohen RA, Gonin R, et al. Lipid levels in the second year of life among HIV-infected and HIV-exposed uninfected Latin American children. *AIDS*. 2012;26(2):235-240. Available at <http://www.ncbi.nlm.nih.gov/pubmed/22008654>.
26. Arpadi S, Shiao S, Strehlau R, et al. Metabolic abnormalities and body composition of HIV-infected children on Lopinavir or Nevirapine-based antiretroviral therapy. *Arch Dis Child*. 2013;98(4):258-264. Available at <http://www.ncbi.nlm.nih.gov/pubmed/23220209>.
27. Calvo M, Martinez E. Update on metabolic issues in HIV patients. *Curr Opin HIV AIDS*. 2014;9(4):332-339. Available at <http://www.ncbi.nlm.nih.gov/pubmed/24824886>.
28. Barlow-Mosha L, Eckard AR, McComsey GA, Musoke PM. Metabolic complications and treatment of perinatally HIV-infected children and adolescents. *J Int AIDS Soc*. 2013;16:18600. Available at <http://www.ncbi.nlm.nih.gov/pubmed/23782481>.
29. Blazquez D, Ramos-Amador JT, Sainz T, et al. Lipid and glucose alterations in perinatally-acquired HIV-infected adolescents and young adults. *BMC Infect Dis*. 2015;15:119. Available at <http://www.ncbi.nlm.nih.gov/pubmed/25880777>.
30. Bwakura-Dangarembizi M, Musiime V, Szubert AJ, et al. Prevalence of lipodystrophy and metabolic abnormalities in HIV-infected African children after 3 years on first-line antiretroviral therapy. *Pediatr Infect Dis J*. 2015;34(2):e23-31. Available at <http://www.ncbi.nlm.nih.gov/pubmed/25068287>.

31. Dejckhamron P, Unachak K, Aupibul L, Sirisanthana V. Insulin resistance and lipid profiles in HIV-infected Thai children receiving lopinavir/ritonavir-based highly active antiretroviral therapy. *J Pediatr Endocrinol Metab*. 2014;27(5-6):403-412. Available at <http://www.ncbi.nlm.nih.gov/pubmed/24259240>.
32. Ramteke S, Shiao S, Foca M, et al. Growth and lipid profiles in a south african cohort of HIV+ children and HIV- controls. Presented at: 22nd Conference on Retroviruses and Opportunistic Infections. 2015. Seattle, WA.
33. Innes S, Abdullah KL, Haubrich R, Cotton MF, Browne SH. High Prevalence of Dyslipidemia and Insulin Resistance in HIV-Infected Pre-Pubertal African Children on Antiretroviral Therapy. *Pediatr Infect Dis J*. 2015. Available at <http://www.ncbi.nlm.nih.gov/pubmed/26421804>.
33. Tauber WB, Lewis LL. Clinical Review of Elvitegravir/Cobicistat/Emtricitabine/Tenofovir alafenamide (Genvoya). 2015. Available at <http://www.fda.gov/downloads/drugs/developmentapprovalprocess/developmentresources/ucm478088.pdf>.
35. Sax PE, Wohl D, Yin MT, et al. Tenofovir alafenamide versus tenofovir disoproxil fumarate, coformulated with elvitegravir, cobicistat, and emtricitabine, for initial treatment of HIV-1 infection: two randomised, double-blind, phase 3, non-inferiority trials. *Lancet*. 2015;385(9987):2606-2615. Available at <http://www.ncbi.nlm.nih.gov/pubmed/25890673>.

Table 12c. Antiretroviral-Therapy-Associated Adverse Effects and Management Recommendations—Gastrointestinal Effects (Last updated March 1, 2016; last reviewed March 1, 2016)

Adverse Effects	Associated ARVs	Onset/Clinical Manifestations	Estimated Frequency	Risk Factors	Prevention/Monitoring	Management
Nausea/Vomiting	Principally ZDV and PIs (e.g., LPV/r, RTV), but can occur with all ARVs and COBI	<u>Onset:</u> • Early <u>Presentation:</u> • Nausea, emesis—may be associated with anorexia and/or abdominal pain.	Varies with ARV agent; 10% to 30% in some series	Unknown	Instruct patient to take PIs with food. Generally improves with time; monitor for weight loss, ARV adherence.	Reassure patient/caretaker that nausea and vomiting will likely decrease over time. Provide supportive care, including instruction on dietary modification. Although antiemetics are not generally indicated, they may be useful in extreme or persistent cases.
Diarrhea	PIs (particularly NFV, LPV/r, FPV/r), buffered ddI, INSTI (mild)	<u>Onset:</u> • Early <u>Presentation:</u> • Generally soft, more frequent stools	Varies with ARV agent; 10% to 30% in some series	Unknown	Generally improves with time (usually over 6–8 weeks); monitor for weight loss, dehydration.	Exclude infectious causes of diarrhea. Although data in children on treatment of ARV-associated diarrhea are lacking, dietary modification, use of calcium carbonate (should not be used with DTG), bulk-forming agents (psyllium), or antimotility agents (loperamide) may be helpful. While there are few published data on its use, crofelemer is FDA-approved for treatment of ART-associated diarrhea in adults but not in children.
Pancreatitis	ddI, d4T (especially concurrently or with TDF), boosted PIs Reported, albeit rarely, with most ARVs.	<u>Onset:</u> • Any time, usually after months of therapy <u>Presentation:</u> • Emesis, abdominal pain, elevated amylase and lipase (asymptomatic hyperamylasemia or elevated lipase do not in and of themselves indicate pancreatitis).	<2% in recent series Frequency was higher in the past with higher dosing of ddI.	Concomitant treatment with other medications associated with pancreatitis (e.g., TMP-SMX, pentamidine, ribavirin) Hypertriglyceridemia Advanced disease Previous episode of pancreatitis	Avoid use of ddI in patients with a history of pancreatitis.	Discontinue offending agent—avoid reintroduction. Manage symptoms of acute episode. If associated with hypertriglyceridemia, consider interventions to lower TG levels.

Key to Acronyms: ART = antiretroviral therapy; ARV = antiretroviral; **COBI = cobicistat**; d4T = stavudine; ddI = didanosine; DTG = dolutegravir; FDA = Food and Drug Administration; FPV/r = fosamprenavir/ritonavir; **INSTI = integrase strand transfer inhibitor**; LPV/r = lopinavir/ritonavir; NFV = nelfinavir; PI = protease inhibitor; RTV = ritonavir; TDF = tenofovir disoproxil fumarate; TG = triglyceride; TMP-SMX = trimethoprim sulfamethoxazole; ZDV = zidovudine

References

1. Buck WC, Kabue MM, Kazembe PN, Kline MW. Discontinuation of standard first-line antiretroviral therapy in a cohort of 1434 Malawian children. *J Int AIDS Soc*. 2010;13:31. Available at <http://www.ncbi.nlm.nih.gov/pubmed/20691049>.
2. Kumarasamy N, Venkatesh KK, Devaleenol B, Poongulali S, Mothi SN, Solomon S. Safety, tolerability and effectiveness of generic HAART in HIV-infected children in South India. *J Trop Pediatr*. 2009;55(3):155-159. Available at <http://www.ncbi.nlm.nih.gov/pubmed/18829638>.
3. Nachman SA, Chernoff M, Gona P, et al. Incidence of noninfectious conditions in perinatally HIV-infected children and adolescents in the HAART era. *Arch Pediatr Adolesc Med*. 2009;163(2):164-171. Available at <http://www.ncbi.nlm.nih.gov/pubmed/19188649>.
4. Hoffmann CJ, Fielding KL, Charalambous S, et al. Antiretroviral therapy using zidovudine, lamivudine, and efavirenz in South Africa: tolerability and clinical events. *AIDS*. 2008;22(1):67-74. Available at <http://www.ncbi.nlm.nih.gov/pubmed/18090393>.
5. Malan N, Su J, Mancini M, et al. Gastrointestinal tolerability and quality of life in antiretroviral-naïve HIV-1-infected patients: data from the CASTLE study. *AIDS Care*. 2010;22(6):677-686. Available at <http://www.ncbi.nlm.nih.gov/pubmed/20467943>.
6. Manfredi R, Calza L. HIV infection and the pancreas: risk factors and potential management guidelines. *Int J STD AIDS*. 2008;19(2):99-105. Available at <http://www.ncbi.nlm.nih.gov/pubmed/18334062>.
7. Turner MJ, Angel JB, Woodend K, Giguere P. The efficacy of calcium carbonate in the treatment of protease inhibitor-induced persistent diarrhea in HIV-infected patients. *HIV Clin Trials*. 2004;5(1):19-24. Available at <http://www.ncbi.nlm.nih.gov/pubmed/15002083>.
8. Heiser CR, Ernst JA, Barrett JT, French N, Schutz M, Dube MP. Probiotics, soluble fiber, and L-Glutamine (GLN) reduce nelfinavir (NFV)- or lopinavir/ritonavir (LPV/r)-related diarrhea. *J Int Assoc Physicians AIDS Care (Chic)*. 2004;3(4):121-129. Available at <http://www.ncbi.nlm.nih.gov/pubmed/15768732>.
9. Tukei VJ, Asimwe A, Maganda A, et al. Safety and tolerability of antiretroviral therapy among HIV-infected children and adolescents in Uganda. *J Acquir Immune Defic Syndr*. 2012;59(3):274-280. Available at <http://www.ncbi.nlm.nih.gov/pubmed/22126740>.
10. Wegzyn CM, Fredrick LM, Stubbs RO, Woodward WC, Norton M. Diarrhea Associated with Lopinavir/Ritonavir-Based Therapy: Results of a Meta-Analysis of 1469 HIV-1-Infected Participants. *J Int Assoc Physicians AIDS Care (Chic)*. 2012;11(4):252-259. Available at <http://www.ncbi.nlm.nih.gov/pubmed/22544446>.
11. Oumar AA, Diallo K, Dembele JP, et al. Adverse drug reactions to antiretroviral therapy: prospective study in children in sikasso (mali). *J Pediatr Pharmacol Ther*. 2012;17(4):382-388. Available at <http://www.ncbi.nlm.nih.gov/pubmed/23411444>.
12. Patel TS, Crutchley RD, Tucker AM, Cottreau J, Garey KW. Crofelemer for the treatment of chronic diarrhea in patients living with HIV/AIDS. *HIV/AIDS*. 2013;5:153-162. Available at <http://www.ncbi.nlm.nih.gov/pubmed/23888120>.
13. Wattanuchariya N, Sirisanthana V, Oberdorfer P. Effectiveness and safety of protease inhibitor-based regimens in HIV-infected Thai children failing first-line treatment. *HIV Med*. 2013;14(4):226-232. Available at <http://www.ncbi.nlm.nih.gov/pubmed/23094820>.
14. Van Dyke RB, Wang L, Williams PL, Pediatric ACTGCT. Toxicities associated with dual nucleoside reverse-transcriptase inhibitor regimens in HIV-infected children. *J Infect Dis*. 2008;198(11):1599-1608. Available at <http://www.ncbi.nlm.nih.gov/pubmed/19000014>.
15. Strehlau R, Donati AP, Arce PM, et al. PRINCE-1: safety and efficacy of atazanavir powder and ritonavir liquid in HIV-1-infected antiretroviral-naïve and -experienced infants and children aged ≥ 3 months to < 6 years. *J Int AIDS Soc*. 2015;18:19467. Available at <http://www.ncbi.nlm.nih.gov/pubmed/26066346>.
16. Clay PG, Crutchley RD. Noninfectious Diarrhea in HIV Seropositive individuals: a review of prevalence rates, etiology, and management in the era of combination antiretroviral therapy. *Infect Dis Ther*. 2014. Available at <http://www.ncbi.nlm.nih.gov/pubmed/25388760>.
17. Gutierrez Mdel M, Mateo MG, Vidal F, Domingo P. Drug safety profile of integrase strand transfer inhibitors. *Expert Opin Drug Saf*. 2014;13(4):431-445. Available at <http://www.ncbi.nlm.nih.gov/pubmed/24597519>.
18. Lee FJ, Carr A. Tolerability of HIV integrase inhibitors. *Curr Opin HIV AIDS*. 2012;7(5):422-428. Available at <http://www.ncbi.nlm.nih.gov/pubmed/22886031>.

Table 12d. Antiretroviral-Therapy-Associated Adverse Effects and Management Recommendations—Hematologic Effects (Last updated March 1, 2016; last reviewed March 1, 2016) (page 1 of 2)

Adverse Effects	Associated ARVs	Onset/Clinical Manifestations	Estimated Frequency	Risk Factors	Prevention/Monitoring	Management
Anemia^a	Principally ZDV	<p><u>Onset:</u></p> <ul style="list-style-type: none"> • Variable, weeks to months <p><u>Presentation</u></p> <p><i>Most Commonly:</i></p> <ul style="list-style-type: none"> • Asymptomatic or mild fatigue • Pallor • Tachypnea <p><i>Rarely:</i></p> <ul style="list-style-type: none"> • Congestive heart failure 	<p><u>HIV-Exposed Newborns:</u></p> <ul style="list-style-type: none"> • Severe anemia is uncommon, but may be seen coincident with physiologic Hgb nadir. <p><u>HIV-Infected Children on ARVs:</u></p> <ul style="list-style-type: none"> • 2–3 times more common with ZDV-containing regimens; less frequent with currently recommended dosing of ZDV 	<p><u>HIV-Exposed Newborns:</u></p> <ul style="list-style-type: none"> • Premature birth • <i>In utero</i> exposure to ARVs • Advanced maternal HIV • Neonatal blood loss • Combination ARV prophylaxis, particularly with ZDV plus 3TC <p><u>HIV-Infected Children on ARVs:</u></p> <ul style="list-style-type: none"> • Underlying hemoglobinopathy (e.g., sickle cell disease, G6PD deficiency) • Myelosuppressive drugs (e.g., TMP-SMX, rifabutin) • Iron deficiency • Advanced or poorly controlled HIV disease • Malnutrition 	<p><u>HIV-Exposed Newborns:</u></p> <ul style="list-style-type: none"> • Obtain CBC at birth. • Consider repeat CBC at 4 weeks for neonates who are at higher risk (e.g., those born prematurely or known to have low birth Hgb). <p><u>HIV-Infected Children on ARVs:</u></p> <ul style="list-style-type: none"> • Avoid ZDV in children with moderate to severe anemia when alternative agents are available. • Obtain CBC as part of routine care. 	<p><u>HIV-Exposed Newborns:</u></p> <ul style="list-style-type: none"> • Rarely require intervention unless Hgb is <7.0 g/dL or anemia is associated with symptoms. • Consider discontinuing ZDV if 4 weeks or more of a 6-week ZDV prophylaxis regimen are already completed (see the Perinatal Guidelines^b). <p><u>HIV-Infected Children on ARVs:</u></p> <ul style="list-style-type: none"> • Discontinue non-ARV, marrow-toxic drugs, if feasible. • Treat coexisting iron deficiency, OIs, malignancies. • For persistent severe anemia thought to be associated with ARVs, change to a non-ZDV-containing regimen; consider a trial of erythropoietin if essential to continue ZDV.
Macrocytosis	Principally ZDV; also d4T	<p><u>Onset:</u></p> <ul style="list-style-type: none"> • Within days to weeks of starting therapy • MCV often >100 fL <p><u>Presentation:</u></p> <ul style="list-style-type: none"> • Most often asymptomatic • Sometimes associated with anemia (occurs more often with ZDV than with d4T) 	>90% to 95%, all ages	None	Obtain CBC as part of routine care.	None required unless associated with anemia

Table 12d. Antiretroviral-Therapy-Associated Adverse Effects and Management Recommendations—Hematologic Effects (Last updated March 1, 2016; last reviewed March 1, 2016) (page 2 of 2)

Adverse Effects	Associated ARVs	Onset/Clinical Manifestations	Estimated Frequency	Risk Factors	Prevention/Monitoring	Management
Neutropenia^a	Principally ZDV	<p><u>Onset:</u></p> <ul style="list-style-type: none"> • Variable <p><u>Presentation:</u></p> <ul style="list-style-type: none"> • Most commonly asymptomatic. Complications appear to be less than with neutropenias associated with cancer chemotherapy. 	<p><u>HIV-Exposed Newborns:</u></p> <ul style="list-style-type: none"> • Rare <p><u>HIV-Infected Children on ARVs:</u></p> <ul style="list-style-type: none"> • 2.2% to 26.8% of children on ARVs, depending upon the ARV regimen. 2.2% for ZDV/3TC • Highest rates with ZDV-containing regimens 	<p><u>HIV-Exposed Newborns:</u></p> <ul style="list-style-type: none"> • <i>In utero</i> exposure to ARVs • Combination ARV prophylaxis, particularly with ZDV plus 3TC <p><u>HIV-Infected Children on ARVs:</u></p> <ul style="list-style-type: none"> • Advanced or poorly controlled HIV infection • Myelosuppressive drugs (e.g., TMP-SMX, ganciclovir, hydroxyurea, rifabutin) 	<p><u>HIV-Infected Children on ARVs:</u></p> <ul style="list-style-type: none"> • Obtain CBC as part of routine care. 	<p><u>HIV-Exposed Newborns:</u></p> <ul style="list-style-type: none"> • No established threshold for intervention; some experts would consider using an alternative NRTI for prophylaxis if ANC <500 cells/mm³, or discontinue ARV prophylaxis entirely if ≥4 weeks of 6-week ZDV prophylaxis have been completed (see the Perinatal Guidelines^b). <p><u>HIV-Infected Children on ARVs:</u></p> <ul style="list-style-type: none"> • Discontinue non-ARV marrow-toxic drugs, if feasible. • Treat coexisting OIs and malignancies. • For persistent severe neutropenia thought to be associated with ARVs, change to a non-ZDV-containing regimen. Consider a trial of G-CSF if essential to continue ZDV.

^a HIV infection itself, OIs, and medications used to prevent OIs, such as TMP-SMX, may all contribute to anemia, neutropenia, and thrombocytopenia.

^b *Recommendations for Use of Antiretroviral Drugs in Pregnant HIV-1-Infected Women for Maternal Health and Interventions to Reduce Perinatal HIV Transmission in the United States*

Key to Acronyms: 3TC = lamivudine; ANC = absolute neutrophil count; ARV = antiretroviral; CBC = complete blood count; d4t = stavudine; dL = deciliter; fL = femtoliter; G6PD = glucose-6-phosphate dehydrogenase; G-CSF = granulocyte colony-stimulating factor; Hgb = hemoglobin; MCV = mean cell volume; NRTI = nucleoside reverse transcriptase inhibitor; OI = opportunistic infection; TMP-SMX = trimethoprim-sulfamethoxazole; ZDV = zidovudine

References

1. Englund JA, Baker CJ, Raskino C, et al. Zidovudine, didanosine, or both as the initial treatment for symptomatic HIV-infected children. AIDS Clinical Trials Group (ACTG) Study 152 Team. *N Engl J Med*. 1997;336(24):1704-1712. Available at <http://www.ncbi.nlm.nih.gov/pubmed/9182213>.
2. Starr SE, Fletcher CV, Spector SA, et al. Combination therapy with efavirenz, nelfinavir, and nucleoside reverse-transcriptase inhibitors in children infected with human immunodeficiency virus type 1. Pediatric AIDS Clinical Trials Group 382 Team. *N Engl J Med*. 1999;341(25):1874-1881. Available at <http://www.ncbi.nlm.nih.gov/pubmed/10601506>.
3. Connor EM, Sperling RS, Gelber R, et al. Reduction of maternal-infant transmission of human immunodeficiency virus type 1 with zidovudine treatment. Pediatric AIDS Clinical Trials Group Protocol 076 Study Group. *N Engl J Med*. 1994;331(18):1173-1180. Available at <http://www.ncbi.nlm.nih.gov/pubmed/7935654>.
4. Krogstad P, Lee S, Johnson G, et al. Nucleoside-analogue reverse-transcriptase inhibitors plus nevirapine, nelfinavir, or ritonavir for pretreated children infected with human immunodeficiency virus type 1. *Clin Infect Dis*. 2002;34(7):991-1001. Available at <http://www.ncbi.nlm.nih.gov/pubmed/11880966>.
5. Najean Y, Rain JD. The mechanism of thrombocytopenia in patients with HIV infection. *J Lab Clin Med*. 1994;123(3):415-420. Available at <http://www.ncbi.nlm.nih.gov/pubmed/8133154>.
6. Caselli D, Maccabruni A, Zuccotti GV, et al. Recombinant erythropoietin for treatment of anaemia in HIV-infected children. *AIDS*. 1996;10(8):929-931. Available at <http://www.ncbi.nlm.nih.gov/pubmed/8828757>.
7. Allen UD, Kirby MA, Goeree R. Cost-effectiveness of recombinant human erythropoietin versus transfusions in the treatment of zidovudine-related anemia in HIV-infected children. *Pediatr AIDS HIV Infect*. 1997;8(1):4-11. Available at <http://www.ncbi.nlm.nih.gov/pubmed/11361510>.
8. Mueller BU, Jacobsen F, Butler KM, Husson RN, Lewis LL, Pizzo PA. Combination treatment with azidothymidine and granulocyte colony-stimulating factor in children with human immunodeficiency virus infection. *J Pediatr*. 1992;121(5 Pt 1):797-802. Available at <http://www.ncbi.nlm.nih.gov/pubmed/1279153>.
9. Scaradavou A, Woo B, Woloski BM, et al. Intravenous anti-D treatment of immune thrombocytopenic purpura: experience in 272 patients. *Blood*. 1997;89(8):2689-2700. Available at <http://www.ncbi.nlm.nih.gov/pubmed/9108386>.
10. Lahoz R, Noguera A, Rovira N, et al. Antiretroviral-related hematologic short-term toxicity in healthy infants: implications of the new neonatal 4-week zidovudine regimen. *Pediatr Infect Dis J*. 2010;29(4):376-379. Available at <http://www.ncbi.nlm.nih.gov/pubmed/19949355>.
11. Dryden-Peterson S, Shapiro RL, Hughes MD, et al. Increased risk of severe infant anemia after exposure to maternal HAART, Botswana. *J Acquir Immune Defic Syndr*. 2011;56(5):428-436. Available at <http://www.ncbi.nlm.nih.gov/pubmed/21266910>.
12. Mocroft A, Lifson AR, Touloumi G, et al. Haemoglobin and anaemia in the SMART study. *Antivir Ther*. 2011;16(3):329-337. Available at <http://www.ncbi.nlm.nih.gov/pubmed/21555815>.
13. Nyesigire Ruhinda E, Bajunirwe F, Kiwanuka J. Anaemia in HIV-infected children: severity, types and effect on response to HAART. *BMC Pediatr*. 2012;12:170. Available at <http://www.ncbi.nlm.nih.gov/pubmed/23114115>.
14. Esan MO, Jonker FA, Hensbroek MB, Calis JC, Phiri KS. Iron deficiency in children with HIV-associated anaemia: a systematic review and meta-analysis. *Trans R Soc Trop Med Hyg*. 2012;106(10):579-587. Available at <http://www.ncbi.nlm.nih.gov/pubmed/22846115>.
15. Nielsen-Saines K, Watts DH, Veloso VG, et al. Three postpartum antiretroviral regimens to prevent intrapartum HIV infection. *N Engl J Med*. 2012;366(25):2368-2379. Available at <http://www.ncbi.nlm.nih.gov/pubmed/22716975>.
16. Renner LA, Dicko F, Koueta F, et al. Anaemia and zidovudine-containing antiretroviral therapy in paediatric antiretroviral programmes in the IeDEA Paediatric West African Database to evaluate AIDS. *J Int AIDS Soc*. 2013;16(1):18024. Available at <http://www.ncbi.nlm.nih.gov/pubmed/24047928>.
17. Arrow Trial team, Kekitiinwa A, Cook A, et al. Routine versus clinically driven laboratory monitoring and first-line antiretroviral therapy strategies in African

children with HIV (ARROW): a 5-year open-label randomised factorial trial. *Lancet*. 2013;381(9875):1391-1403. Available at <http://www.ncbi.nlm.nih.gov/pubmed/23473847>.

18. Bunupuradah T, Kariminia A, Chan KC, et al. Incidence and predictors of severe anemia in Asian HIV-infected children using first-line antiretroviral therapy. *Int J Infect Dis*. 2013;17(10):e806-810. Available at <http://www.ncbi.nlm.nih.gov/pubmed/23764352>.
19. Singh A, Hemal A, Agarwal S, Dubey N, Buxi G. A prospective study of haematological changes after switching from stavudine to zidovudine-based antiretroviral treatment in HIV-infected children. *Int J STD AIDS*. 2014. Available at <http://www.ncbi.nlm.nih.gov/pubmed/24516076>.
20. Van Dyke RB, Wang L, Williams PL, Pediatric ACTGCT. Toxicities associated with dual nucleoside reverse-transcriptase inhibitor regimens in HIV-infected children. *J Infect Dis*. 2008;198(11):1599-1608. Available at <http://www.ncbi.nlm.nih.gov/pubmed/19000014>.
21. Smith C, Forster JE, Levin MJ, et al. Serious adverse events are uncommon with combination neonatal antiretroviral prophylaxis: a retrospective case review. *PLoS One*. 2015;10(5):e0127062. Available at <http://www.ncbi.nlm.nih.gov/pubmed/26000984>.

Table 12e. Antiretroviral-Therapy-Associated Adverse Effects and Management Recommendations—Hepatic Events

(Last updated March 1, 2016; last reviewed March 1, 2016) (page 1 of 2)

Adverse Effects	Associated ARVs	Onset/Clinical Manifestations	Estimated Frequency	Risk Factors	Prevention/Monitoring	Management
Hepatic Toxicity Elevated AST, ALT, clinical hepatitis	<p>All ARVs may be associated with hepatitis. NVP and TPV are of particular concern.</p> <p>NVP, EFV, ABC, RAL, and MVC have been associated with hypersensitivity reactions.</p> <p>NRTIs (especially ZDV, ddI, and d4T) are associated with lactic acidosis and hepatic steatosis.</p>	<p><u>Onset:</u></p> <ul style="list-style-type: none"> Hepatitis generally occurs within the first few months of therapy, but can occur later. Steatosis presents after months to years of therapy. HBV-coinfected patients may develop severe hepatic flare with the initiation, withdrawal, or development of resistance to 3TC, FTC, or TDF (especially in patients receiving only one anti-HBV agent). Hepatitis may also represent IRIS early in therapy, especially in HBV- and HCV-infected patients. <p><u>Presentation:</u></p> <ul style="list-style-type: none"> Asymptomatic elevation of AST and ALT Symptomatic hepatitis with nausea, fatigue, and jaundice Hepatitis may be component of hypersensitivity reaction with rash, lactic acidosis, and hepatic steatosis. 	<p>Uncommon in children</p> <p>Frequency varies with different agents and drug combinations.</p>	<p>HBV or HCV coinfection</p> <p>Elevated baseline ALT and AST</p> <p>Other hepatotoxic medications (including herbal preparations such as St. John's wort [<i>Hypericum perforatum</i>], Chaparral [<i>Larrea tridentate</i>], German chamomile [<i>Matricaria inodora</i>], and others)</p> <p>Alcohol use</p> <p>Underlying liver disease</p> <p>Pregnancy</p> <p><u>For NVP-Associated Hepatic Events in Adults:</u></p> <ul style="list-style-type: none"> Female with pre-NVP CD4 count >250 cells/mm³ Male with pre-NVP CD4 count >400 cells/mm³ <p>Certain HLA types are also associated with NVP-associated hepatic events but are population-specific.^a Higher drug concentrations for PIs, particularly TPV.</p>	<p><u>Prevention:</u></p> <ul style="list-style-type: none"> Avoid concomitant use of hepatotoxic medications. If hepatic enzymes are elevated >5 to 10 times ULN or chronic liver disease, most clinicians would avoid NVP. <p><u>Monitoring:</u></p> <p><i>For ARVs Other Than NVP:</i></p> <ul style="list-style-type: none"> Obtain AST and ALT at baseline and thereafter at least every 3–4 months, or more frequently in at-risk patients (e.g., HBV- or HCV-coinfected or elevated baseline AST and ALT). <p><i>For NVP:</i></p> <ul style="list-style-type: none"> Obtain AST and ALT at baseline, at 2 and 4 weeks, then every 3 months. 	<p>Asymptomatic patients with elevated ALT or AST should be evaluated for other causes and monitored closely</p> <p>(including repeating AST, ALT and checking total bilirubin). If ALT or AST is more than 5–10 times ULN and felt to be possibly or probably associated with ARVs, the potentially offending ARVs should be discontinued.</p> <p>In symptomatic patients, discontinue all ARVs and other potential hepatotoxic agents and avoid restarting the offending agent.</p> <p>If a symptomatic hepatic event occurs on NVP, permanently discontinue drug (see also NVP Hypersensitivity).</p> <p>When clinical hepatitis is associated with lactic acidosis, avoid restarting the most likely agent, including ZDV, d4T, and ddI in particular (see also Lactic Acidosis).</p> <p>Consider viral causes of hepatitis: HAV, HBV, HCV, EBV, and CMV.</p>

Table 12e. Antiretroviral-Therapy-Associated Adverse Effects and Management Recommendations—Hepatic Events
(Last updated March 1, 2016; last reviewed March 1, 2016) (page 2 of 2)

Adverse Effects	Associated ARVs	Onset/Clinical Manifestations	Estimated Frequency	Risk Factors	Prevention/Monitoring	Management
Indirect Hyperbilirubinemia	IDV, ATV (with either RTV or COBI)	<p><u>Onset:</u></p> <ul style="list-style-type: none"> First months of therapy <p><u>Presentation:</u></p> <ul style="list-style-type: none"> Jaundice; otherwise asymptomatic elevation of indirect bilirubin levels with normal AST, and ALT. <p>Direct bilirubin may be normal or slightly elevated when levels of indirect bilirubin are very high.</p>	<p><u>HIV-Infected Children Receiving ATV:</u></p> <ul style="list-style-type: none"> In long-term follow-up, 9% had at least 1 total bilirubin level > 5 x ULN and 1.4% experienced jaundice 	N/A	<p><u>Monitoring:</u></p> <ul style="list-style-type: none"> No specific monitoring. 	<p>Not necessary to discontinue the offending agent except for cosmetic reasons.</p> <p>After an initial rise over the first few months of therapy, unconjugated bilirubin levels generally stabilize; in some patients, levels improve over time.</p>
Non-Cirrhotic Portal Hypertension	ddl, d4t	<p><u>Onset:</u></p> <ul style="list-style-type: none"> Generally after years of therapy <p><u>Presentation:</u></p> <ul style="list-style-type: none"> GI bleeding, esophageal varices, hypersplenism Mild elevations in AST and ALT, moderate increases in ALP, and pancytopenia (because of hypersplenism) Liver biopsy may reveal a variety of findings, most commonly nodular regenerative hyperplasia or hepatoportal sclerosis. 	<p><u>Rare:</u></p> <ul style="list-style-type: none"> Probably less than 1% 	Prolonged exposure to ARV therapy, especially ddl and the combination of ddl and d4T	<p><u>Monitoring:</u></p> <ul style="list-style-type: none"> No specific monitoring 	<p>Manage complications of GI bleeding and esophageal varices.</p> <p>Discontinue/replace d4T or ddl, if patient is receiving either.</p>

^a For example, HLA-DRB1*0101 in whites, HLA-DRB1*0102 in South Africans, and HLA-B35 in Thai and whites.

Key to Acronyms: 3TC = lamivudine; ABC = abacavir; ALP = alkaline phosphatase; ALT = alanine transaminase; ARV = antiretroviral; AST = aspartate aminotransferase; ATV = atazanavir; CD4 = CD4 T lymphocyte; CMV = cytomegalovirus; **COBI = cobicistat**; d4T = stavudine; ddl = didanosine; EBV = Epstein-Barr virus; EFV = efavirenz; FTC = emtricitabine; GI = gastrointestinal; HAV = hepatitis A virus; HBV = hepatitis B virus; HCV = hepatitis C virus; HLA = human leukocyte antigen; IDV = indinavir; IRIS = immune reconstitution inflammatory syndrome; MVC = maraviroc; NRTI = nucleoside reverse transcriptase inhibitor; NVP = nevirapine; PI = protease inhibitor; RAL = raltegravir; **RTV = ritonavir**; TDF = tenofovir disoproxil fumarate; TPV = tipranavir; ULN = upper limit of normal; ZDV = zidovudine

References

1. Aceti A, Pasquazzi C, Zechini B, De Bac C, Group L. Hepatotoxicity development during antiretroviral therapy containing protease inhibitors in patients with HIV: the role of hepatitis B and C virus infection. *J Acquir Immune Defic Syndr*. 2002;29(1):41-48. Available at <http://www.ncbi.nlm.nih.gov/pubmed/11782588>.
2. Baylor MS, Johann-Liang R. Hepatotoxicity associated with nevirapine use. *J Acquir Immune Defic Syndr*. 2004;35(5):538-539. Available at http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=pubmed&dopt=Abstract&list_uids=15021321.
3. Buck WC, Kabue MM, Kazembe PN, Kline MW. Discontinuation of standard first-line antiretroviral therapy in a cohort of 1434 Malawian children. *J Int AIDS Soc*. 2010;13:31. Available at <http://www.ncbi.nlm.nih.gov/pubmed/20691049>.
4. Bunchorntavakul C, Reddy KR. Review article: herbal and dietary supplement hepatotoxicity. *Aliment Pharmacol Ther*. 2013;37(1):3-17. Available at <http://www.ncbi.nlm.nih.gov/pubmed/23121117>.
5. Busti AJ, Hall RG, Margolis DM. Atazanavir for the treatment of human immunodeficiency virus infection. *Pharmacotherapy*. 2004;24(12):1732-1747. Available at <http://www.ncbi.nlm.nih.gov/pubmed/15585441>.
6. Cotte L, Benet T, Billioud C, et al. The role of nucleoside and nucleotide analogues in nodular regenerative hyperplasia in HIV-infected patients: a case control study. *J Hepatol*. 2011;54(3):489-496. Available at <http://www.ncbi.nlm.nih.gov/pubmed/21056493>.
7. Gray D, Nuttall J, Lombard C, et al. Low rates of hepatotoxicity in HIV-infected children on anti-retroviral therapy with and without isoniazid prophylaxis. *Journal of tropical pediatrics*. 2010;56(3):159-165. Available at <http://www.ncbi.nlm.nih.gov/pubmed/19710246>.
8. Kovari H, Ledergerber B, Battegay M, et al. Incidence and risk factors for chronic elevation of alanine aminotransferase levels in HIV-infected persons without hepatitis b or c virus co-infection. *Clin Infect Dis*. 2010;50(4):502-511. Available at <http://www.ncbi.nlm.nih.gov/pubmed/20085465>.
9. Kovari H, Ledergerber B, Peter U, et al. Association of noncirrhotic portal hypertension in HIV-infected persons and antiretroviral therapy with didanosine: a nested case-control study. *Clin Infect Dis*. 2009;49(4):626-635. Available at <http://www.ncbi.nlm.nih.gov/pubmed/19589079>.
10. Levy V, Grant RM. Antiretroviral therapy for hepatitis B virus-HIV-coinfected patients: promises and pitfalls. *Clin Infect Dis*. 2006;43(7):904-910. Available at <http://www.ncbi.nlm.nih.gov/pubmed/16941375>.
11. McDonald C, Uy J, Hu W, et al. Clinical significance of hyperbilirubinemia among HIV-1-infected patients treated with atazanavir/ritonavir through 96 weeks in the CASTLE study. *AIDS Patient Care STDS*. 2012;26(5):259-264. Available at <http://www.ncbi.nlm.nih.gov/pubmed/22404426>.
12. McKoy JM, Bennett CL, Scheetz MH, et al. Hepatotoxicity associated with long- versus short-course HIV-prophylactic nevirapine use: a systematic review and meta-analysis from the Research on Adverse Drug events And Reports (RADAR) project. *Drug safety: an international journal of medical toxicology and drug experience*. 2009;32(2):147-158. Available at <http://www.ncbi.nlm.nih.gov/pubmed/19236121>.
13. Nunez M. Clinical syndromes and consequences of antiretroviral-related hepatotoxicity. *Hepatology*. 2010;52(3):1143-1155. Available at <http://www.ncbi.nlm.nih.gov/pubmed/20812358>.
14. Ouyang DW, Shapiro DE, Lu M, et al. Increased risk of hepatotoxicity in HIV-infected pregnant women receiving antiretroviral therapy independent of nevirapine exposure. *AIDS*. 2009;23(18):2425-2430. Available at <http://www.ncbi.nlm.nih.gov/pubmed/19617813>.
15. Phillips E, Bartlett JA, Sanne I, et al. Associations between HLA-DRB1*0102, HLA-B*5801, and hepatotoxicity during initiation of nevirapine-containing regimens in South Africa. *J Acquir Immune Defic Syndr*. 2013;62(2):e55-57. Available at <http://www.ncbi.nlm.nih.gov/pubmed/23328091>.
16. Schouten JN, Van der Ende ME, Koeter T, et al. Risk factors and outcome of HIV-associated idiopathic noncirrhotic portal hypertension. *Aliment Pharmacol Ther*. 2012;36(9):875-885. Available at <http://www.ncbi.nlm.nih.gov/pubmed/22971050>.

17. Stern JO, Robinson PA, Love J, Lanes S, Imperiale MS, Mayers DL. A comprehensive hepatic safety analysis of nevirapine in different populations of HIV infected patients. *J Acquir Immune Defic Syndr*. 2003;34 Suppl 1(Suppl 1):S21-33. Available at <http://www.ncbi.nlm.nih.gov/pubmed/14562855>.
18. Van Dyke RB, Wang L, Williams PL, Pediatric ACTGCT. Toxicities associated with dual nucleoside reverse-transcriptase inhibitor regimens in HIV-infected children. *J Infect Dis*. 2008;198(11):1599-1608. Available at <http://www.ncbi.nlm.nih.gov/pubmed/19000014>.
19. Vispo E, Morello J, Rodriguez-Novoa S, Soriano V. Noncirrhotic portal hypertension in HIV infection. *Curr Opin Infect Dis*. 2011;24(1):12-18. Available at <http://www.ncbi.nlm.nih.gov/pubmed/21157331>.
20. Wit FW, Weverling GJ, Weel J, Jurriaans S, Lange JM. Incidence of and risk factors for severe hepatotoxicity associated with antiretroviral combination therapy. *J Infect Dis*. 2002;186(1):23-31. Available at <http://www.ncbi.nlm.nih.gov/pubmed/12089658>.
21. Aupibul L, Bunupuradah T, Sophan S, et al. Prevalence and incidence of liver dysfunction and assessment of biomarkers of liver disease in HIV-infected Asian children. *Pediatr Infect Dis J*. 2015;34(6):e153-158. Available at <http://www.ncbi.nlm.nih.gov/pubmed/25970117>.
22. Strehlau R, Donati AP, Arce PM, et al. PRINCE-1: safety and efficacy of atazanavir powder and ritonavir liquid in HIV-1-infected antiretroviral-naïve and -experienced infants and children aged ≥ 3 months to < 6 years. *J Int AIDS Soc*. 2015;18:19467. Available at <http://www.ncbi.nlm.nih.gov/pubmed/26066346>.
23. Rutstein RM, Samson, P., Fenton, T., Fletcher, C.V., Kiser, J.J., Mofenson, L.M., Smith, E., Graham, B., Mathew, M., Aldrovani, G.; for the PACTG 1020A Study Team. Long-Term Safety and Efficacy of Atazanavir-Based Therapy in HIV-Infected Infants, Children and Adolescents: The Pediatric AIDS Clinical Trials Group Protocol 1020A. *Pediatr Infect Dis J*. 2015;34:162-167. Available at
24. Food and Drug Administration. TYBOST (Cobicistat) [package insert]. Available at http://www.gilead.com/~media/Files/pdfs/medicines/hiv/tybost/tybost_pi.pdf. 2014.

Table 12f. Antiretroviral-Therapy-Associated Adverse Effects and Management Recommendations—Insulin Resistance, Asymptomatic Hyperglycemia, Diabetes Mellitus (Last updated March 1, 2016; last reviewed March 1, 2016)

Adverse Effects	Associated ARVs	Onset/Clinical Manifestations	Estimated Frequency	Risk Factors	Prevention/Monitoring	Management
Insulin Resistance, Asymptomatic Hyperglycemia, DM^a	Several NRTIs (e.g., d4T, ZDV, ddI) Several PIs (e.g., LPV/r; less often ATV, ATV/r, DRV/r, NFV, TPV/r)	<u>Onset:</u> • Weeks to months after beginning therapy; median of 60 days (adult data). <u>Presentation</u> <i>Most Commonly:</i> • Asymptomatic fasting hyperglycemia (possibly in the setting of lipodystrophy), metabolic syndrome, or growth delay <i>Also Possible:</i> • Frank DM (i.e., polyuria, polydipsia, polyphagia, fatigue, hyperglycemia)	<u>Insulin Resistance</u> <i>ARV-Treated Adults and Children:</i> • 6% to 33% <u>Impaired Fasting Glucose</u> <i>ARV-Treated Adults:</i> • 3% to 25% <i>ARV-Treated Children:</i> • 0% to 7% <u>Impaired Glucose Tolerance</u> <i>ARV-Treated Adults:</i> • 16% to 35% <i>ARV-Treated Children:</i> • 3% to 4% <u>DM</u> <i>ARV-Treated Adults:</i> • 0.6–4.7 per 100 person-years (2- to 4-fold greater than that for HIV-uninfected adults) <i>ARV-Treated Children:</i> • Rare in HIV-infected children	<u>Risk Factors for Type 2 DM:</u> • Lipodystrophy • Metabolic syndrome • Family history of DM • High BMI (obesity)	<u>Prevention:</u> • Lifestyle modification • Although uncertain, avoiding the use of d4T may reduce risk. <u>Monitoring:</u> • Monitor for polydipsia, polyuria, polyphagia, change in body habitus, and acanthosis nigricans. <i>Obtain RPG Levels at:</i> • Initiation of ARV therapy • 3–6 months after therapy initiation • Once a year thereafter <i>For RPG ≥ 140 mg/dL:</i> • Obtain FPG performed after 8-hour fast and consider referral to endocrinologist.	Counsel on lifestyle modification (e.g., a diet low in saturated fat, cholesterol, trans fat, and refined sugars; increased physical activity; cessation of smoking); consider consultation with dietician. Change NRTI (e.g., from d4T, ZDV, or ddI to TDF or ABC). <u>For Either RPG ≥ 200 mg/dL Plus Symptoms of DM or FPG ≥ 126 mg/dL:</u> • Patient meets diagnostic criteria for DM; consult endocrinologist. <u>FPG 100–125 mg/dL:</u> • Impaired FPG is suggestive of insulin resistance; consult endocrinologist <u>FPG < 100 mg/dL:</u> <i>Normal FPG, but Does Not Exclude Insulin Resistance:</i> • Recheck FPG in 6–12 months.

^a Insulin resistance, asymptomatic hyperglycemia, and DM form a spectrum of increasing severity. *Insulin resistance* is often defined as elevated insulin levels for the level of glucose observed; *impaired FPG* as an FPG of 100–125 mg/dL; *impaired glucose tolerance* as an elevated 2-hour PG of 140–199 mg/dL in a 75g-OGTT (or if <43 kg, 1.75 g/kg of glucose up to a maximum of 75g); and *diabetes mellitus* as either an FPG ≥ 126 mg/dL, a random PG ≥ 200 mg/dL in a patient with hyperglycemia symptoms, an HgbA1C of ≥ 6.5%, or a 2-hour PG after OGTT ≥ 200 mg/dL. However, the Panel does not recommend routine determinations of insulin levels, HgbA1C, or glucose tolerance without consultation with an endocrinologist; these guidelines are instead based on the readily available random and fasting plasma glucose levels.

Key to Acronyms: ABC = abacavir; ARV = antiretroviral; ATV = atazanavir; ATV/r = ritonavir-boosted atazanavir; BMI = body mass index; d4T = stavudine; ddI = didanosine; **dL = deciliter**; DM = diabetes mellitus; DRV/r = ritonavir-boosted darunavir; FPG = fasting plasma glucose; HgbA1c = glycosylated hemoglobin; LPV/r = ritonavir-boosted lopinavir; NFV = nelfinavir; NRTI = nucleoside reverse transcriptase inhibitor; OGTT = oral glucose tolerance test; PG = plasma glucose; PI = protease inhibitor; RPG = random plasma glucose; TDF = tenofovir disoproxil fumarate; TPV/r = ritonavir-boosted tipranavir; ZDV = zidovudine

References

1. Bitnun A, Sochetti E, Dick PT, et al. Insulin sensitivity and beta-cell function in protease inhibitor-treated and -naive human immunodeficiency virus-infected children. *J Clin Endocrinol Metab*. 2005;90(1):168-174. Available at <http://www.ncbi.nlm.nih.gov/pubmed/15483082>.
2. Hadigan C. Insulin resistance among HIV-infected patients: unraveling the mechanism. *Clin Infect Dis*. 2005;41(9):1341-1342. Available at <http://www.ncbi.nlm.nih.gov/pubmed/16206113>.
3. Morse CG, Kovacs JA. Metabolic and skeletal complications of HIV infection: the price of success. *JAMA*. 2006;296(7):844-854. Available at <http://www.ncbi.nlm.nih.gov/pubmed/16905789>.
4. Brown TT, Cole SR, Li X, et al. Antiretroviral therapy and the prevalence and incidence of diabetes mellitus in the multicenter AIDS cohort study. *Arch Intern Med*. 2005;165(10):1179-1184. Available at <http://www.ncbi.nlm.nih.gov/pubmed/15911733>.
5. Justman JE, Benning L, Danoff A, et al. Protease inhibitor use and the incidence of diabetes mellitus in a large cohort of HIV-infected women. *J Acquir Immune Defic Syndr*. 2003;32(3):298-302. Available at <http://www.ncbi.nlm.nih.gov/pubmed/12626890>.
6. Aldrovandi GM, Lindsey JC, Jacobson DL, et al. Morphologic and metabolic abnormalities in vertically HIV-infected children and youth. *AIDS*. 2009;23(6):661-672. Available at <http://www.ncbi.nlm.nih.gov/pubmed/19279441>.
7. Chantry CJ, Hughes MD, Alvero C, et al. Lipid and glucose alterations in HIV-infected children beginning or changing antiretroviral therapy. *Pediatrics*. 2008;122(1):e129-138. Available at <http://www.ncbi.nlm.nih.gov/pubmed/18519448>.
8. Samaras K. Prevalence and pathogenesis of diabetes mellitus in HIV-1 infection treated with combined antiretroviral therapy. *J Acquir Immune Defic Syndr*. 2009;50(5):499-505. Available at <http://www.ncbi.nlm.nih.gov/pubmed/19223782>.
9. Geffner ME, Patel K, Miller TL, et al. Factors associated with insulin resistance among children and adolescents perinatally infected with HIV-1 in the pediatric HIV/AIDS cohort study. *Horm Res Paediatr*. 2011;76(6):386-391. Available at <http://www.ncbi.nlm.nih.gov/pubmed/22042056>.
10. Petoumenos K, Worm SW, Fontas E, et al. Predicting the short-term risk of diabetes in HIV-positive patients: the Data Collection on Adverse Events of Anti-HIV Drugs (D:A:D) study. *J Int AIDS Soc*. 2012;15(2):17426. Available at <http://www.ncbi.nlm.nih.gov/pubmed/23078769>.
11. Rasmussen LD, Mathiesen ER, Kronborg G, Pedersen C, Gerstoft J, Obel N. Risk of diabetes mellitus in persons with and without HIV: a Danish nationwide population-based cohort study. *PLoS One*. 2012;7(9):e44575. Available at <http://www.ncbi.nlm.nih.gov/pubmed/22984529>.
12. American Diabetes Association. Standards of medical care in diabetes-2015. *Diabetes care*. 2015;38 Suppl:S1-2. Available at <http://www.ncbi.nlm.nih.gov/pubmed/25537700>.
13. Nathan DM, Buse JB, Davidson MB, et al. Medical management of hyperglycemia in type 2 diabetes: a consensus algorithm for the initiation and adjustment of therapy: a consensus statement of the American Diabetes Association and the European Association for the Study of Diabetes. *Diabetes Care*. 2009;32(1):193-203. Available at <http://www.ncbi.nlm.nih.gov/pubmed/18945920>.
14. Nathan DM, Davidson MB, DeFronzo RA, et al. Impaired fasting glucose and impaired glucose tolerance: implications for care. *Diabetes Care*. 2007;30(3):753-759. Available at <http://www.ncbi.nlm.nih.gov/pubmed/17327355>.
15. Feeney ER, Mallon PW. Insulin resistance in treated HIV infection. *Best Pract Res Clin Endocrinol Metab*. 2011;25(3):443-458. Available at <http://www.ncbi.nlm.nih.gov/pubmed/21663838>.
16. Hazra R, Hance LF, Monteiro JP, et al. Insulin resistance and glucose and lipid concentrations in a cohort of perinatally HIV-infected Latin American children. *Pediatr Infect Dis J*. 2013;32(7):757-759. Available at <http://www.ncbi.nlm.nih.gov/pubmed/23360832>.

17. Patel K, Wang J, Jacobson DL, et al. Aggregate risk of cardiovascular disease among adolescents perinatally infected with the human immunodeficiency virus. *Circulation*. 2014;129(11):1204-1212. Available at <http://www.ncbi.nlm.nih.gov/pubmed/24366631>.
18. Hadigan C, Kattakuzhy S. Diabetes mellitus type 2 and abnormal glucose metabolism in the setting of human immunodeficiency virus. *Endocrinol Metab Clin North Am*. 2014;43(3):685-696. Available at <http://www.ncbi.nlm.nih.gov/pubmed/25169561>.
19. Fortuny C, Deya-Martinez A, Chiappini E, Galli L, de Martino M, Noguera-Julian A. Metabolic and renal adverse effects of antiretroviral therapy in HIV-infected children and adolescents. *Pediatr Infect Dis J*. 2015;34(5 Suppl 1):S36-43. Available at <http://www.ncbi.nlm.nih.gov/pubmed/25629891>.
20. Loomba-Albrecht LA, Bregman T, Chantry CJ. Endocrinopathies in children infected with human immunodeficiency virus. *Endocrinol Metab Clin North Am*. 2014;43(3):807-828. Available at <http://www.ncbi.nlm.nih.gov/pubmed/25169569>.
21. Innes S, Abdullah KL, Haubrich R, Cotton MF, Browne SH. High Prevalence of Dyslipidemia and Insulin Resistance in HIV-Infected Pre-Pubertal African Children on Antiretroviral Therapy. *Pediatr Infect Dis J*. 2015. Available at <http://www.ncbi.nlm.nih.gov/pubmed/26421804>.

Table 12g. Antiretroviral-Therapy-Associated Adverse Effects and Management Recommendations—Lactic Acidosis

(Last updated March 1, 2016; last reviewed March 1, 2016)

Adverse Effects	Associated ARVs	Onset/Clinical Manifestations	Estimated Frequency	Risk Factors	Prevention/Monitoring	Management
Lactic Acidosis	NRTIs, in particular, d4T and ddI (highest risk when co-administered)	<p>Onset:</p> <ul style="list-style-type: none"> 1–20 months after starting therapy (median onset 4 months in 1 case series) <p>Presentation</p> <p><i>Usually Insidious Onset of a Combination of Signs and Symptoms:</i></p> <ul style="list-style-type: none"> Generalized fatigue, weakness, and myalgias Vague abdominal pain, weight loss, unexplained nausea or vomiting Dyspnea Peripheral neuropathy <p>Note: Patients may present with acute multi-organ failure (e.g., fulminant hepatic, pancreatic, respiratory failure).</p>	<p>Chronic, Asymptomatic Mild Hyperlactatemia (2.1–5.0 mmol/L)</p> <p><i>Adults:</i></p> <ul style="list-style-type: none"> 15% to 35% of adults receiving NRTI therapy for longer than 6 months <p><i>Children:</i></p> <ul style="list-style-type: none"> 29% to 32% <p>Symptomatic Severe Hyperlactatemia (>5.0 mmol/L)</p> <p><i>Adults:</i></p> <ul style="list-style-type: none"> 0.2% to 5.7% <p>Symptomatic Lactic Acidosis/Hepatic Steatosis:</p> <ul style="list-style-type: none"> Rare in all age groups (1.3–11 episodes per 1000 person-years; increased incidence with the use of d4T/ddI when co-administered), but associated with a high fatality rate (33% to 58%) 	<p>Adults:</p> <ul style="list-style-type: none"> Female gender High BMI Chronic HCV infection African-American race Prolonged NRTI use (particularly d4T and ddI) Co-administration of ddI with other agents (e.g., d4T, TDF, RBV, tetracycline) Co-administration of TDF with metformin Overdose of propylene glycol CD4 count <350 cells/mm³ Acquired riboflavin or thiamine deficiency Possibly pregnancy <p>Preterm Infants:</p> <ul style="list-style-type: none"> Exposure to propylene glycol (e.g., present as a diluent in LPV/r oral solution) 	<p>Prevention:</p> <ul style="list-style-type: none"> Avoid d4T and ddI individually; co-administration of d4T and ddI is not recommended in an ARV regimen (no exception). Due to the presence of propylene glycol as a diluent, LPV/r oral solution should never be used in preterm neonates in the immediate postnatal period. Monitor for clinical manifestations of lactic acidosis and promptly adjust therapy. <p>Monitoring</p> <p><i>Asymptomatic:</i></p> <ul style="list-style-type: none"> Measurement of serum lactate is not recommended. <p><i>Clinical Signs or Symptoms Consistent with Lactic Acidosis:</i></p> <ul style="list-style-type: none"> Obtain blood lactate level.^a Additional diagnostic evaluations should include serum bicarbonate and anion gap and/or arterial blood gas, amylase and lipase, serum albumin, and hepatic transaminases. 	<p>Lactate 2.1–5.0 mmol/L (Confirmed with Second Test):</p> <ul style="list-style-type: none"> Consider replacing ddI and d4T with other ARVs. As an alternative, temporarily discontinue all ARVs while conducting additional diagnostic workup. <p>Lactate >5.0 mmol/L (Confirmed with Second Test)^b or >10.0 mmol/L (Any 1 Test):</p> <ul style="list-style-type: none"> Discontinue all ARVs. Provide supportive therapy (IV fluids; some patients may require sedation and respiratory support to reduce oxygen demand and ensure adequate oxygenation of tissues). <p>Anecdotal (Unproven) Supportive Therapies:</p> <ul style="list-style-type: none"> Bicarbonate infusions, THAM, high-dose thiamine and riboflavin, oral antioxidants (e.g., L-carnitine, co-enzyme Q10, vitamin C) <p>Following resolution of clinical and laboratory abnormalities, resume therapy, either with an NRTI-sparing regimen or a revised NRTI-containing regimen instituted with caution, using NRTIs less likely to inhibit mitochondria (ABC or TDF preferred; possibly FTC or 3TC), and monthly monitoring of lactate for at least 3 months.</p>

^a Blood for lactate determination should be collected, without prolonged tourniquet application or fist clenching, into a pre-chilled, gray-top, fluoride-oxalate-containing tube and transported on ice to the laboratory to be processed within 4 hours of collection.

^b Management can be initiated before the results of the confirmatory test.

Key to Acronyms: 3TC = lamivudine; ABC = abacavir; ARV = antiretroviral; BMI = body mass index; CD4 = CD4 T lymphocyte; d4T = stavudine; ddI = didanosine; FTC = emtricitabine; HCV = hepatitis C virus; IV = intravenous; LPV/r = lopinavir/ritonavir; NRTI = nucleoside reverse transcriptase inhibitor; RBV = ribavirin; TDF = tenofovir disoproxil fumarate; THAM = tris (hydroxymethyl) aminomethane

References

General Reviews

1. Fortuny C, Deya-Martinez A, Chiappini E, Galli L, de Martino M, Noguera-Julian A. Metabolic and renal adverse effects of antiretroviral therapy in HIV-infected children and adolescents. *Pediatr Infect Dis J*. 2015;34(5 Suppl 1):S36-43. Available at <http://www.ncbi.nlm.nih.gov/pubmed/25629891>.
2. Arenas-Pinto A, Grant A, Bhaskaran K, et al. Risk factors for fatality in HIV-infected patients with dideoxynucleoside-induced severe hyperlactataemia or lactic acidosis. *Antivir Ther*. 2011;16(2):219-226. Available at <http://www.ncbi.nlm.nih.gov/pubmed/21447871>.
3. Tukei VJ, Asimwe A, Maganda A, et al. Safety and tolerability of antiretroviral therapy among HIV-infected children and adolescents in Uganda. *J Acquir Immune Defic Syndr*. 2012;59(3):274-280. Available at <http://www.ncbi.nlm.nih.gov/pubmed/22126740>.

Risk Factors

4. Manosuthi W, Prasithsirikul W, Chumpathat N, et al. Risk factors for mortality in symptomatic hyperlactatemia among HIV-infected patients receiving antiretroviral therapy in a resource-limited setting. *Int J Infect Dis*. 2008;12(6):582-586. Available at <http://www.ncbi.nlm.nih.gov/pubmed/18337140>.
5. Osler M, Stead D, Rebe K, Meintjes G, Boulle A. Risk factors for and clinical characteristics of severe hyperlactataemia in patients receiving antiretroviral therapy: a case-control study. *HIV Med*. 2010;11(2):121-129. Available at <http://www.ncbi.nlm.nih.gov/pubmed/19702629>.
6. Aperis G, Paliouras C, Zervos A, Arvanitis A, Alivanis P. Lactic acidosis after concomitant treatment with metformin and tenofovir in a patient with HIV infection. *Journal of Renal Care*. 2011;37(1):25-29. Available at <http://www.ncbi.nlm.nih.gov/pubmed/21288314>.
7. Boxwell DC, K.; et al. Neonatal Toxicity of Kaletra Oral Solution—LPV, Ethanol, or Propylene Glycol?— Abstract #708. 18th Conference on Retroviruses and Opportunistic Infections (CROI). Boston MA. 2011. Available at
8. Feeney ER, Chazallon C, O'Brien N, et al. Hyperlactataemia in HIV-infected subjects initiating antiretroviral therapy in a large randomized study (a substudy of the INITIO trial). *HIV Med*. 2011;12(10):602-609. Available at <http://www.ncbi.nlm.nih.gov/pubmed/21599820>.
9. Leung L, Wilson D, Manini AF. Fatal toxicity from symptomatic hyperlactataemia: a retrospective cohort study of factors implicated with long-term nucleoside reverse transcriptase inhibitor use in a South African hospital. *Drug Saf*. 2011;34(6):521-527. Available at <http://www.ncbi.nlm.nih.gov/pubmed/21488705>.
10. Maskew M, Westreich D, Fox MP, Maotse T, Sanne IM. Effectiveness and safety of 30 mg versus 40 mg stavudine regimens: a cohort study among HIV-infected adults initiating HAART in South Africa. *J Int AIDS Soc*. 2012;15(1):13. Available at <http://www.ncbi.nlm.nih.gov/pubmed/22410312>.
11. Matthews LT, Giddy J, Ghebremichael M, et al. A risk-factor guided approach to reducing lactic acidosis and hyperlactatemia in patients on antiretroviral therapy. *PLoS One*. 2011;6(4):e18736. Available at <http://www.ncbi.nlm.nih.gov/pubmed/21494566>.
12. Menezes CN, Maskew M, Sanne I, Crowther NJ, Raal FJ. A longitudinal study of stavudine-associated toxicities in a large cohort of South African HIV infected subjects. *BMC Infect Dis*. 2011;11:244. Available at <http://www.ncbi.nlm.nih.gov/pubmed/21923929>.
13. Phan V, Thai S, Choun K, Lynen L, van Griensven J. Incidence of treatment-limiting toxicity with stavudine-based antiretroviral therapy in Cambodia: a retrospective cohort study. *PLoS One*. 2012;7(1):e30647. Available at <http://www.ncbi.nlm.nih.gov/pubmed/22303447>.
14. Dragovic G, Jevtovic D. The role of nucleoside reverse transcriptase inhibitors usage in the incidence of hyperlactatemia and lactic acidosis in HIV/AIDS patients. *Biomedicine & pharmacotherapy = Biomedecine & pharmacotherapie*. 2012;66(4):308-311. Available at <http://www.ncbi.nlm.nih.gov/pubmed/22658063>.
15. Moren C, Noguera-Julian A, Garrabou G, et al. Mitochondrial evolution in HIV-infected children receiving first- or second-generation nucleoside analogues. *J Acquir Immune Defic Syndr*. 2012;60(2):111-116. Available at <http://www.ncbi.nlm.nih.gov/pubmed/22362155>.
16. Palmer M, Chersich M, Moultrie H, Kuhn L, Fairlie L, Meyers T. Frequency of stavudine substitution due to toxicity in children receiving antiretroviral treatment in

sub-Saharan Africa. *AIDS*. 2013;27(5):781-785. Available at <http://www.ncbi.nlm.nih.gov/pubmed/23169331>.

17. Wester CW, Eden SK, Shepherd BE, et al. Risk factors for symptomatic hyperlactatemia and lactic acidosis among combination antiretroviral therapy-treated adults in Botswana: results from a clinical trial. *AIDS Res Hum Retroviruses*. 2012;28(8):759-765. Available at <http://www.ncbi.nlm.nih.gov/pubmed/22540188>.
18. Mamiafo CT, Moor VJ, Nansseu JR, Pieme CA, Tayou C, Yonkeu JN. Hyperlactatemia in a group of HIV patients living in Yaounde-Cameroon. *AIDS Res Ther*. 2014;11(1):2. Available at <http://www.ncbi.nlm.nih.gov/pubmed/24428886>.
19. Margolis AM, Heverling H, Pham PA, Stolbach A. A review of the toxicity of HIV medications. *Journal of medical toxicology: official journal of the American College of Medical Toxicology*. 2014;10(1):26-39. Available at <http://www.ncbi.nlm.nih.gov/pubmed/23963694>.

Monitoring and Management

20. Barlow-Mosha L, Eckard AR, McComsey GA, Musoke PM. Metabolic complications and treatment of perinatally HIV-infected children and adolescents. *J Int AIDS Soc*. 2013;16:18600. Available at <http://www.ncbi.nlm.nih.gov/pubmed/23782481>.
21. Kraut JA, Madias NE. Lactic acidosis. *N Engl J Med*. 2014;371(24):2309-2319. Available at <http://www.ncbi.nlm.nih.gov/pubmed/25494270>.
22. Claessens YE, Cariou A, Monchi M, et al. Detecting life-threatening lactic acidosis related to nucleoside-analog treatment of human immunodeficiency virus-infected patients, and treatment with L-carnitine. *Critical care medicine*. 2003;31(4):1042-1047. Available at <http://www.ncbi.nlm.nih.gov/pubmed/12682470>.
23. Marfo K, Garala M, Kvetan V, Gasperino J. Use of Tris-hydroxymethyl aminomethane in severe lactic acidosis due to highly active antiretroviral therapy: a case report. *Journal of clinical pharmacy and therapeutics*. 2009;34(1):119-123. Available at <http://www.ncbi.nlm.nih.gov/pubmed/19125910>.
24. McComsey G, Loneragan JT. Mitochondrial dysfunction: patient monitoring and toxicity management. *J Acquir Immune Defic Syndr*. 2004;37 Suppl 1:S30-35. Available at <http://www.ncbi.nlm.nih.gov/pubmed/15319667>.
25. Chagoma N, Mallewa J, Kaunda S, et al. Longitudinal lactate levels from routine point-of-care monitoring in adult Malawian antiretroviral therapy patients: associations with stavudine toxicities. *Trans R Soc Trop Med Hyg*. 2013;107(10):615-619. Available at <http://www.ncbi.nlm.nih.gov/pubmed/23926161>.

Table 12h. Antiretroviral-Therapy-Associated Adverse Effects and Management Recommendations—Lipodystrophy, Lipohypertrophy, Lipoatrophy (Last updated March 1, 2016; last reviewed March 1, 2016) (page 1 of 2)

Adverse Effects	Associated ARVs	Onset/Clinical Manifestations	Estimated Frequency	Risk Factors	Prevention/Monitoring	Management
Lipodystrophy (Fat Maldistribution) General Information	See below for specific associations.	<u>Onset:</u> • Trunk and limb fat initially increase within a few months of start of ART; peripheral fat wasting may not appear for 12 to 24 months after ART initiation.	Varies greatly depending upon measure and comparator group <u>Highly Variable in Adults:</u> • Up to 93% <u>Children:</u> • Up to 34%, perhaps more common in adolescents than prepubertal children	Genetic predisposition Puberty HIV-associated inflammation Older age Longer duration of ART Body habitus	See below.	See below Although more typically associated with certain ARVs (e.g., d4T), a regimen review with consideration of changing the regimen should be considered, whenever present
Central Lipohypertrophy or Lipo-accumulation	Can occur in the absence of ART, but most associated with PIs and EFV.	<u>Presentation:</u> • Central fat accumulation with increased abdominal girth, which may include dorsocervical fat pad (buffalo hump) and/or gynecomastia in males or breast hypertrophy in females, particularly with EFV . The appearance of central lipohypertrophy is accentuated in the presence of peripheral fat wasting (lipoatrophy).	<u>Adults:</u> • Up to 93% <u>Children:</u> • Up to 27%	Obesity before initiation of therapy Sedentary lifestyle	<u>Prevention:</u> • Calorically appropriate low-fat diet and exercise <u>Monitoring:</u> • BMI measurement • Body circumference and waist-hip ratio	Calorically appropriate healthy diet low in saturated fats and simple carbohydrates , and exercise, especially strength training Smoking cessation (if applicable) to decrease future CVD risk Consider switching from PIs and EFV to an INSTI <u>Data are Insufficient to Allow the Panel to Safely Recommend Use of Any of the Following Modalities in Children:</u> • Recombinant human growth hormone • Growth hormone-releasing hormone • Metformin • Thiazolidinediones • Anabolic steroids • Liposuction.

Table 12h. Antiretroviral-Therapy-Associated Adverse Effects and Management Recommendations—Lipodystrophy, Lipohypertrophy, Lipoatrophy (Last updated March 1, 2016; last reviewed March 1, 2016) (page 2 of 2)

Adverse Effects	Associated ARVs	Onset/Clinical Manifestations	Estimated Frequency	Risk Factors	Prevention/Monitoring	Management
Facial/Peripheral Lipoatrophy	Most associated with thymidine analogue NRTIs (d4T > ZDV)	<u>Presentation:</u> <ul style="list-style-type: none"> Thinning of subcutaneous fat in face, buttocks, and extremities, measured as decrease in trunk/limb fat by DXA or triceps skinfold thickness. Preservation of lean body mass distinguishes lipoatrophy from HIV-associated wasting. 	<u>Adults:</u> <ul style="list-style-type: none"> Up to 59% (particularly in patients on d4T-containing regimens) <u>Children:</u> <ul style="list-style-type: none"> Up to 47% (particularly in patients on d4T-containing regimens) Risk lower (up to 15%) in patients not treated with d4T or ZDV. 	Underweight before ART	<u>Prevention:</u> <ul style="list-style-type: none"> Avoid use of d4T and ZDV. <u>Monitoring:</u> <ul style="list-style-type: none"> Patient self-report and physical exam are the most sensitive methods of monitoring lipoatrophy. 	Replace d4T (not widely used and recommended only in special circumstances) or ZDV with other NRTIs if possible without loss of virologic control. <u>Data are Insufficient to Allow the Panel to Safely Recommend Use of Any of the Following Modalities in Children:</u> <ul style="list-style-type: none"> Injections of poly-L-lactic acid Recombinant human leptin Autologous fat transplantation Thiazolidinediones.

Key to Acronyms: ART = antiretroviral therapy; ARV = antiretroviral; BMI = body mass index; CVD = cardiovascular disease; d4T = stavudine; DXA = dual energy x-ray absorptiometry; EFV = efavirenz; NRTI = nucleoside reverse transcriptase inhibitor; PI = protease inhibitor; ZDV = zidovudine

References

See the archived version of Supplement III, February 23, 2009 *Guidelines for the Use of Antiretroviral Agents in Pediatric HIV Infection*, (<http://www.aidsinfo.nih.gov>) for a more complete discussion and reference list.

General Reviews

- Fernandez JR, Redden DT, Pietrobelli A, Allison DB. Waist circumference percentiles in nationally representative samples of African-American, European-American, and Mexican-American children and adolescents. *J Pediatr*. 2004;145(4):439-444. Available at <http://www.ncbi.nlm.nih.gov/pubmed/15480363>.
- Lee JM, Davis MM, Woolford SJ, Gurney JG. Waist circumference percentile thresholds for identifying adolescents with insulin resistance in clinical practice. *Pediatric diabetes*. 2009;10(5):336-342. Available at <http://www.ncbi.nlm.nih.gov/pubmed/19175894>.
- Carr A. Treatment strategies for HIV lipodystrophy. *Curr Opin HIV AIDS*. 2007;2(4):332-338. Available at <http://www.ncbi.nlm.nih.gov/pubmed/19372908>.
- Phillips DR, Hay P. Current perspectives on the management and prevention of antiretroviral-associated lipoatrophy. *J Antimicrob Chemother*. 2008;62(5):866-871. Available at <http://www.ncbi.nlm.nih.gov/pubmed/18703527>.
- Moyle G, Moutschen M, Martinez E, et al. Epidemiology, assessment, and management of excess abdominal fat in persons with HIV infection. *AIDS reviews*. 2010;12(1):3-14. Available at <http://www.ncbi.nlm.nih.gov/pubmed/20216906>.
- Dzwonek A, Clapson M, Withey S, Bates A, Novelli V. Severe gynecomastia in an African boy with perinatally acquired human immunodeficiency virus infection receiving highly active antiretroviral therapy. *Pediatr Infect Dis J*. 2006;25(2):183-184. Available at <http://www.ncbi.nlm.nih.gov/pubmed/16462304>.

7. Innes S, Cotton MF, Haubrich R, et al. High prevalence of lipoatrophy in pre-pubertal South African children on antiretroviral therapy: a cross-sectional study. *BMC Pediatr.* 2012;12:183. Available at <http://www.ncbi.nlm.nih.gov/pubmed/23176441>.
8. Piloya T, Bakeera-Kitaka S, Kekitiinwa A, Kamya MR. Lipodystrophy among HIV-infected children and adolescents on highly active antiretroviral therapy in Uganda: a cross sectional study. *J Int AIDS Soc.* 2012;15(2):17427. Available at <http://www.ncbi.nlm.nih.gov/pubmed/22814353>.
9. Prendergast AJ. Complications of long-term antiretroviral therapy in HIV-infected children. *Arch Dis Child.* 2013;98(4):245-246. Available at <http://www.ncbi.nlm.nih.gov/pubmed/23413313>.
10. Arbeitman LE, O'Brien RC, Somarriba G, et al. Body mass index and waist circumference of HIV-infected youth in a Miami cohort: comparison to local and national cohorts. *Journal of pediatric gastroenterology and nutrition.* 2014;59(4):449-454. Available at <http://www.ncbi.nlm.nih.gov/pubmed/24709829>.
11. Bwakura-Dangarembizi M, Musiime V, Szubert AJ, et al. Prevalence of lipodystrophy and metabolic abnormalities in HIV-infected African children after 3 years on first-line antiretroviral therapy. *Pediatr Infect Dis J.* 2015;34(2):e23-31. Available at <http://www.ncbi.nlm.nih.gov/pubmed/25068287>.

Associated ARVs/Etiology

12. Dube MP, Komarow L, Mulligan K, et al. Long-term body fat outcomes in antiretroviral-naïve participants randomized to nelfinavir or efavirenz or both plus dual nucleosides. Dual X-ray absorptiometry results from A5005s, a substudy of Adult Clinical Trials Group 384. *J Acquir Immune Defic Syndr.* 2007;45(5):508-514. Available at <http://www.ncbi.nlm.nih.gov/pubmed/17589373>.
13. Haubrich RH, Riddler SA, DiRienzo AG, et al. Metabolic outcomes in a randomized trial of nucleoside, nonnucleoside and protease inhibitor-sparing regimens for initial HIV treatment. *AIDS.* 2009;23(9):1109-1118. Available at <http://www.ncbi.nlm.nih.gov/pubmed/19417580>.
14. Hulgán T, Tebas P, Canter JA, et al. Hemochromatosis gene polymorphisms, mitochondrial haplogroups, and peripheral lipoatrophy during antiretroviral therapy. *J Infect Dis.* 2008;197(6):858-866. Available at <http://www.ncbi.nlm.nih.gov/pubmed/18419350>.
15. McComsey GA, Libutti DE, O'Riordan M, et al. Mitochondrial RNA and DNA alterations in HIV lipoatrophy are linked to antiretroviral therapy and not to HIV infection. *Antivir Ther.* 2008;13(5):715-722. Available at <http://www.ncbi.nlm.nih.gov/pubmed/18771055>.
16. Tien PC, Benson C, Zolopa AR, Sidney S, Osmond D, Grunfeld C. The study of fat redistribution and metabolic change in HIV infection (FRAM): methods, design, and sample characteristics. *American journal of epidemiology.* 2006;163(9):860-869. Available at <http://www.ncbi.nlm.nih.gov/pubmed/16524955>.
17. Van Dyke RB, Wang L, Williams PL, Pediatric ACTGCT. Toxicities associated with dual nucleoside reverse-transcriptase inhibitor regimens in HIV-infected children. *J Infect Dis.* 2008;198(11):1599-1608. Available at <http://www.ncbi.nlm.nih.gov/pubmed/19000014>.
18. Mulligan K, Parker RA, Komarow L, et al. Mixed patterns of changes in central and peripheral fat following initiation of antiretroviral therapy in a randomized trial. *J Acquir Immune Defic Syndr.* 2006;41(5):590-597. Available at <http://www.ncbi.nlm.nih.gov/pubmed/16652032>.
19. Scherzer R, Shen W, Bacchetti P, et al. Comparison of dual-energy X-ray absorptiometry and magnetic resonance imaging-measured adipose tissue depots in HIV-infected and control subjects. *The American journal of clinical nutrition.* 2008;88(4):1088-1096. Available at <http://www.ncbi.nlm.nih.gov/pubmed/18842798>.
20. Benn P, Sauret-Jackson V, Cartledge J, et al. Improvements in cheek volume in lipoatrophic individuals switching away from thymidine nucleoside reverse transcriptase inhibitors. *HIV Med.* 2009;10(6):351-355. Available at <http://www.ncbi.nlm.nih.gov/pubmed/19490181>.
21. Arpadí S, Shiao S, Strehlau R, et al. Metabolic abnormalities and body composition of HIV-infected children on Lopinavir or Nevirapine-based antiretroviral therapy. *Arch Dis Child.* 2013;98(4):258-264. Available at <http://www.ncbi.nlm.nih.gov/pubmed/23220209>.
22. Foca M, Wang L, Ramteke R, et al. Changes in mitochondrial enzyme function as a predictor of lipodystrophy. Presented at: 7th International AIDS Society. 2015. Vancouver, Canada.
23. Cohen S, Innes S, Geelen SP, et al. Long-term changes of subcutaneous fat mass in HIV-infected children on antiretroviral therapy: a retrospective analysis of longitudinal data from two pediatric HIV-cohorts. *PLoS One.* 2015;10(7):e0120927. Available at <http://www.ncbi.nlm.nih.gov/pubmed/26148119>.
24. Dos Reis LC, de Carvalho Rondo PH, de Sousa Marques HH, Jose Segri N. Anthropometry and body composition of vertically HIV-infected children and adolescents

under therapy with and without protease inhibitors. *Public Health Nutr.* 2015;18(7):1255-1261. Available at <http://www.ncbi.nlm.nih.gov/pubmed/25115797>.

Management

25. Wohl DA, Brown TT. Management of morphologic changes associated with antiretroviral use in HIV-infected patients. *J Acquir Immune Defic Syndr.* 2008;49 Suppl 2:S93-S100. Available at <http://www.ncbi.nlm.nih.gov/pubmed/18725818>.
26. Carey DL, Baker D, Rogers GD, et al. A randomized, multicenter, open-label study of poly-L-lactic acid for HIV-1 facial lipoatrophy. *J Acquir Immune Defic Syndr.* 2007;46(5):581-589. Available at <http://www.ncbi.nlm.nih.gov/pubmed/18193500>.
27. Cavalcanti RB, Raboud J, Shen S, Kain KC, Cheung A, Walmsley S. A randomized, placebo-controlled trial of rosiglitazone for HIV-related lipoatrophy. *J Infect Dis.* 2007;195(12):1754-1761. Available at <http://www.ncbi.nlm.nih.gov/pubmed/17492590>.
28. Falutz J, Allas S, Blot K, et al. Metabolic effects of a growth hormone-releasing factor in patients with HIV. *N Engl J Med.* 2007;357(23):2359-2370. Available at <http://www.ncbi.nlm.nih.gov/pubmed/18057338>.
29. Gerschenson M, Kim C, Berzins B, et al. Mitochondrial function, morphology and metabolic parameters improve after switching from stavudine to a tenofovir-containing regimen. *J Antimicrob Chemother.* 2009;63(6):1244-1250. Available at <http://www.ncbi.nlm.nih.gov/pubmed/19321503>.
30. Hadigan C. Peroxisome proliferator-activated receptor gamma agonists and the treatment of HIV-associated lipoatrophy: unraveling the molecular mechanism of their shortcomings. *J Infect Dis.* 2008;198(12):1729-1731. Available at <http://www.ncbi.nlm.nih.gov/pubmed/18954262>.
31. Lindegaard B, Hansen T, Hvid T, et al. The effect of strength and endurance training on insulin sensitivity and fat distribution in human immunodeficiency virus-infected patients with lipodystrophy. *The Journal of clinical endocrinology and metabolism.* 2008;93(10):3860-3869. Available at <http://www.ncbi.nlm.nih.gov/pubmed/18628529>.
32. Lo J, You SM, Canavan B, et al. Low-dose physiological growth hormone in patients with HIV and abdominal fat accumulation: a randomized controlled trial. *JAMA.* 2008;300(5):509-519. Available at <http://www.ncbi.nlm.nih.gov/pubmed/18677023>.
33. Mulligan K, Khatami H, Schwarz JM, et al. The effects of recombinant human leptin on visceral fat, dyslipidemia, and insulin resistance in patients with human immunodeficiency virus-associated lipoatrophy and hypoleptinemia. *The Journal of clinical endocrinology and metabolism.* 2009;94(4):1137-1144. Available at <http://www.ncbi.nlm.nih.gov/pubmed/19174500>.
34. Tebas P, Zhang J, Hafner R, et al. Peripheral and visceral fat changes following a treatment switch to a non-thymidine analogue or a nucleoside-sparing regimen in HIV-infected subjects with peripheral lipoatrophy: results of ACTG A5110. *J Antimicrob Chemother.* 2009;63(5):998-1005. Available at <http://www.ncbi.nlm.nih.gov/pubmed/19299471>.
35. Tebas P, Zhang J, Yarasheski K, et al. Switching to a protease inhibitor-containing, nucleoside-sparing regimen (lopinavir/ritonavir plus efavirenz) increases limb fat but raises serum lipid levels: results of a prospective randomized trial (AIDS clinical trial group 5125s). *J Acquir Immune Defic Syndr.* 2007;45(2):193-200. Available at <http://www.ncbi.nlm.nih.gov/pubmed/17527093>.
36. Dollfus C, Blanche S, Trocme N, Funck-Brentano I, Bonnet F, Levan P. Correction of facial lipoatrophy using autologous fat transplants in HIV-infected adolescents. *HIV Med.* 2009;10(5):263-268. Available at <http://www.ncbi.nlm.nih.gov/pubmed/19178590>.
37. Cofrancesco J, Jr., Freedland E, McComsey G. Treatment options for HIV-associated central fat accumulation. *AIDS Patient Care STDS.* 2009;23(1):5-18. Available at <http://www.ncbi.nlm.nih.gov/pubmed/19055407>.
38. Degris E, Delpierre C, Sommet A, et al. Longitudinal study of body composition of 101 HIV men with lipodystrophy: dual-energy X-ray criteria for lipodystrophy evolution. *Journal of clinical densitometry: the official journal of the International Society for Clinical Densitometry.* 2010;13(2):237-244. Available at <http://www.ncbi.nlm.nih.gov/pubmed/20347366>.
39. Falutz J, Mamputu JC, Potvin D, et al. Effects of tesamorelin (TH9507), a growth hormone-releasing factor analog, in human immunodeficiency virus-infected patients with excess abdominal fat: a pooled analysis of two multicenter, double-blind placebo-controlled phase 3 trials with safety extension data. *The Journal of*

clinical endocrinology and metabolism. 2010;95(9):4291-4304. Available at <http://www.ncbi.nlm.nih.gov/pubmed/20554713>.

40. Ferrer E, del Rio L, Martinez E, et al. Impact of switching from lopinavir/ritonavir to atazanavir/ritonavir on body fat redistribution in virologically suppressed HIV-infected adults. *AIDS Res Hum Retroviruses*. 2011;27(10):1061-1065. Available at <http://www.ncbi.nlm.nih.gov/pubmed/21166602>.
41. Negredo E, Miro O, Rodriguez-Santiago B, et al. Improvement of mitochondrial toxicity in patients receiving a nucleoside reverse-transcriptase inhibitor-sparing strategy: results from the Multicenter Study with Nevirapine and Kaletra (MULTINEKA). *Clin Infect Dis*. 2009;49(6):892-900. Available at <http://www.ncbi.nlm.nih.gov/pubmed/19663689>.
42. Raboud JM, Diong C, Carr A, et al. A meta-analysis of six placebo-controlled trials of thiazolidinedione therapy for HIV lipoatrophy. *HIV Clin Trials*. 2010;11(1):39-50. Available at <http://www.ncbi.nlm.nih.gov/pubmed/20400410>.
43. Sheth SH, Larson RJ. The efficacy and safety of insulin-sensitizing drugs in HIV-associated lipodystrophy syndrome: a meta-analysis of randomized trials. *BMC Infect Dis*. 2010;10:183. Available at <http://www.ncbi.nlm.nih.gov/pubmed/20573187>.
44. Tungsiripat M, Bejjani DE, Rizk N, et al. Rosiglitazone improves lipoatrophy in patients receiving thymidine-sparing regimens. *AIDS*. 2010;24(9):1291-1298. Available at <http://www.ncbi.nlm.nih.gov/pubmed/20453626>.
45. Spoulou V, Kanaka-Gantenbein C, Bathrellou I, et al. Monitoring of lipodystrophic and metabolic abnormalities in HIV-1 infected children on antiretroviral therapy. *Hormones*. 2011;10(2):149-155. Available at <http://www.ncbi.nlm.nih.gov/pubmed/21724540>.
46. Minami R, Yamamoto M, Takahama S, Ando H, Miyamura T, Suematsu E. Comparison of the influence of four classes of HIV antiretrovirals on adipogenic differentiation: the minimal effect of raltegravir and atazanavir. *Journal of infection and chemotherapy: official journal of the Japan Society of Chemotherapy*. 2011;17(2):183-188. Available at <http://www.ncbi.nlm.nih.gov/pubmed/20706762>.

Table 12i. Antiretroviral-Therapy-Associated Adverse Effects and Management Recommendations—Nephrotoxic Effects
(Last updated March 1, 2016; last reviewed March 1, 2016) (page 1 of 2)

Adverse Effects	Associated ARVs	Onset/Clinical Manifestations	Estimated Frequency	Risk Factors	Prevention/Monitoring	Management
Urolithiasis/Nephrolithiasis	ATV, IDV Although DRV causes crystalluria, it is not associated with increased nephrolithiasis risk.	<u>Onset:</u> • Weeks to months after starting therapy <u>Clinical Findings:</u> • Crystalluria, hematuria, pyuria, flank pain, sometimes increased creatinine	ATV-related nephrolithiasis occurs in <10%.	In adults, elevated urine pH (>5.7) Unknown in children	<u>Prevention:</u> • Maintain adequate hydration. <u>Monitoring:</u> • Obtain urinalysis at least every 6–12 months.	Provide adequate hydration and pain control; consider using alternative ARV.
Renal Dysfunction	TDF	<u>Onset:</u> • Variable; in adults, weeks to months after initiation of therapy. • Hypophosphatemia appears at a median of 18 months. • Glucosuria may have onset after a year of therapy. <u>Presentation:</u> <i>More Common:</i> • Increased serum creatinine, proteinuria, normoglycemic glucosuria. Hypophosphatemia, usually asymptomatic; may present with bone and muscle pain, weakness. <i>Less Common:</i> • Renal failure, acute tubular necrosis, Fanconi syndrome, proximal renal tubulopathy, interstitial nephritis, nephrogenic diabetes insipidus with polyuria	<u>Adults:</u> • Approximately 2% with increased serum creatinine • Approximately 0.5% with severe renal complications <u>Children:</u> • Approximately 4% with hypophosphatemia or proximal tubulopathy; higher with prolonged TDF therapy, in advanced HIV infection or concomitant use of ddI	<u>Risk May Be Increased in Children:</u> • Aged >6 years • Black race, Hispanic/Latino ethnicity • Advanced HIV infection • Concurrent use of ddI or PIs (especially LPV/r), and preexisting renal dysfunction • Risk increases with longer duration of TDF treatment.	Monitor urine protein and glucose or urinalysis, and serum creatinine at 3- to 6-month intervals. For patients taking TDF, some panelists add serum phosphate to the list of routine labs to monitor. In the presence of persistent proteinuria or glucosuria, or for symptoms of bone pain or muscle pain or weakness, also measure serum phosphate. Because toxicity risk increases with duration of TDF treatment, frequency of monitoring should not decrease with time. While unproven, routine monitoring intervals of every 3–6 months might be considered. Abnormal values should be confirmed by repeat testing, and frequency of monitoring can be increased if abnormalities are found and TDF is continued.	If TDF is the likely cause, consider using alternative ARV.

Table 12i. Antiretroviral-Therapy-Associated Adverse Effects and Management Recommendations—Nephrotoxic Effects

(Last updated March 1, 2016; last reviewed March 1, 2016) (page 2 of 2)

Adverse Effects	Associated ARVs	Onset/Clinical Manifestations	Estimated Frequency	Risk Factors	Prevention/Monitoring	Management
Elevation in Serum Creatinine	DTG, COBI, RPV	<p><u>Onset:</u></p> <ul style="list-style-type: none"> • Within a month of starting treatment <p><u>Presentation:</u></p> <ul style="list-style-type: none"> • Asymptomatic. These drugs decrease renal tubular secretion of creatinine, leading to an increase in measured serum creatinine without a true change in GFR. 	<p>Common</p> <p>Need to distinguish between true change in GFR and other causes.</p> <p>True change might be associated with other medical conditions, continuing rise of serum creatinine with time, and albuminuria.</p>	N/A	<p>Monitor serum creatinine. Assess for renal dysfunction if serum creatinine increases by >0.4 mg/dL or increases are ongoing with time.</p>	<p>No need to change therapy.</p> <p>Reassure patient about the benign nature of the laboratory abnormality.</p>

Key to Acronyms: ARV = antiretroviral; ATV = atazanavir; COBI = cobicistat; ddI = didanosine; DRV = darunavir; DTG = dolutegravir; GFR = glomerular filtration rate; IDV = indinavir; LPV/r = boosted lopinavir/ritonavir; PI = protease inhibitor; RPV = rilpivirine; TDF = tenofovir disoproxil fumarate

References

1. Andiman WA, Chernoff MC, Mitchell C, et al. Incidence of persistent renal dysfunction in human immunodeficiency virus-infected children: associations with the use of antiretrovirals, and other nephrotoxic medications and risk factors. *Pediatr Infect Dis J*. 2009;28(7):619-625. Available at <http://www.ncbi.nlm.nih.gov/pubmed/19561425>.
2. Brennan A, Evans D, Fox M, et al. Renal insufficiency, nephrotoxicity, and mortality among HIV-infected adults on TDF in a South African cohort: a marginal structural models analysis. Presented at: 18th Conference on Retroviruses and Opportunistic Infections. 2011. Boston, MA.
3. Judd A, Boyd KL, Stohr W, et al. Effect of tenofovir disoproxil fumarate on risk of renal abnormality in HIV-1-infected children on antiretroviral therapy: a nested case-control study. *AIDS*. 2010;24(4):525-534. Available at <http://www.ncbi.nlm.nih.gov/pubmed/20139752>.
4. Mueller BU, Nelson RP, Jr., Sleasman J, et al. A phase I/II study of the protease inhibitor ritonavir in children with human immunodeficiency virus infection. *Pediatrics*. 1998;101(3 Pt 1):335-343. Available at <http://www.ncbi.nlm.nih.gov/pubmed/9480994>.
5. Nachman SA, Chernoff M, Gona P, et al. Incidence of noninfectious conditions in perinatally HIV-infected children and adolescents in the HAART era. *Arch Pediatr Adolesc Med*. 2009;163(2):164-171. Available at <http://www.ncbi.nlm.nih.gov/pubmed/19188649>.
6. Riordan A, Judd A, Boyd K, et al. Tenofovir use in human immunodeficiency virus-1-infected children in the United Kingdom and Ireland. *Pediatr Infect Dis J*. 2009;28(3):204-209. Available at <http://www.ncbi.nlm.nih.gov/pubmed/19209091>.
7. Soler-Palacin P, Melendo S, Noguera-Julian A, et al. Prospective study of renal function in HIV-infected pediatric patients receiving tenofovir-containing HAART regimens. *AIDS*. 2011;25(2):171-176. Available at <http://www.ncbi.nlm.nih.gov/pubmed/21076275>.

8. van Rossum AM, Dieleman JP, Fraaij PL, et al. Indinavir-associated asymptomatic nephrolithiasis and renal cortex atrophy in two HIV-1 infected children. *AIDS*. 2001;15(13):1745-1747. Available at <http://www.ncbi.nlm.nih.gov/pubmed/11546957>.
9. van Rossum AM, Dieleman JP, Fraaij PL, et al. Persistent sterile leukocyturia is associated with impaired renal function in human immunodeficiency virus type 1-infected children treated with indinavir. *Pediatrics*. 2002;110(2 Pt 1):e19. Available at <http://www.ncbi.nlm.nih.gov/pubmed/12165618>.
10. Hall AM, Hendry BM, Nitsch D, Connolly JO. Tenofovir-associated kidney toxicity in HIV-infected patients: a review of the evidence. *Am J Kidney Dis*. 2011;57(5):773-780. Available at <http://www.ncbi.nlm.nih.gov/pubmed/21435764>.
11. Herlitz LC, Mohan S, Stokes MB, Radhakrishnan J, D'Agati VD, Markowitz GS. Tenofovir nephrotoxicity: acute tubular necrosis with distinctive clinical, pathological, and mitochondrial abnormalities. *Kidney international*. 2010;78(11):1171-1177. Available at <http://www.ncbi.nlm.nih.gov/pubmed/20811330>.
12. Cooper RD, Wiebe N, Smith N, Keiser P, Naicker S, Tonelli M. Systematic review and meta-analysis: renal safety of tenofovir disoproxil fumarate in HIV-infected patients. *Clin Infect Dis*. 2010;51(5):496-505. Available at <http://www.ncbi.nlm.nih.gov/pubmed/20673002>.
13. Vigano A, Bedogni G, Manfredini V, et al. Long-term renal safety of tenofovir disoproxil fumarate in vertically HIV-infected children, adolescents and young adults: a 60-month follow-up study. *Clin Drug Investig*. 2011;31(6):407-415. Available at <http://www.ncbi.nlm.nih.gov/pubmed/21528939>.
14. Fraaij PL, Verweel G, van Rossum AM, Hartwig NG, Burger DM, de Groot R. Indinavir/low-dose ritonavir containing HAART in HIV-1 infected children has potent antiretroviral activity, but is associated with side effects and frequent discontinuation of treatment. *Infection*. 2007;35(3):186-189. Available at <http://www.ncbi.nlm.nih.gov/pubmed/17565462>.
15. World Health Organization. Technical update on treatment optimization: Use of efavirenz during pregnancy: A public health perspective. 2012. Available at <http://www.who.int/hiv/pub/treatment2/efavirenz/en/>. Accessed February 10, 2016.
16. Purswani M, Patel K, Kopp JB, et al. Tenofovir treatment duration predicts proteinuria in a multiethnic United States Cohort of children and adolescents with perinatal HIV-1 infection. *Pediatr Infect Dis J*. 2013;32(5):495-500. Available at <http://www.ncbi.nlm.nih.gov/pubmed/23249917>.
17. Nishijima T, Hamada Y, Watanabe K, et al. Ritonavir-boosted darunavir is rarely associated with nephrolithiasis compared with ritonavir-boosted atazanavir in HIV-infected patients. *PLoS One*. 2013;8(10):e77268. Available at <http://www.ncbi.nlm.nih.gov/pubmed/24130871>.
18. de Lastours V, Ferrari Rafael De Silva E, Daudon M, et al. High levels of atazanavir and darunavir in urine and crystalluria in asymptomatic patients. *J Antimicrob Chemother*. 2013;68(8):1850-1856. Available at <http://www.ncbi.nlm.nih.gov/pubmed/23599359>.
19. German P, Liu HC, Szwarcberg J, et al. Effect of cobicistat on glomerular filtration rate in subjects with normal and impaired renal function. *J Acquir Immune Defic Syndr*. 2012;61(1):32-40. Available at <http://www.ncbi.nlm.nih.gov/pubmed/22732469>.
20. Yombi JC, Pozniak A, Boffito M, et al. Antiretrovirals and the kidney in current clinical practice: renal pharmacokinetics, alterations of renal function and renal toxicity. *AIDS*. 2014;28(5):621-632. Available at <http://www.ncbi.nlm.nih.gov/pubmed/24983540>.
21. Lim Y, Lyall H, Foster C. Tenofovir-associated nephrotoxicity in children with perinatally-acquired HIV Infection: a single-centre cohort study. *Clin Drug Investig*. 2015;35(5):327-333. Available at <http://www.ncbi.nlm.nih.gov/pubmed/25861908>.
22. Samarawickrama A, Cai M, Smith ER, et al. Simultaneous measurement of urinary albumin and total protein may facilitate decision-making in HIV-infected patients with proteinuria. *HIV Med*. 2012;13(9):526-532. Available at <http://www.ncbi.nlm.nih.gov/pubmed/22413854>.

Table 12j. Antiretroviral-Therapy-Associated Adverse Effects and Management Recommendations—Osteopenia and Osteoporosis (Last updated March 1, 2016; last reviewed March 1, 2016)

Adverse Effects	Associated ARVs	Onset/Clinical Manifestations	Estimated Frequency	Risk Factors	Prevention/Monitoring	Management
Osteopenia and Osteoporosis	Any ART regimen <u>Specific Agents of Possible Concern:</u> • TDF • PIs, especially LPV/r	<u>Onset:</u> • Any age; more common in months after initiation of ART. <u>Presentation:</u> • Most commonly asymptomatic; fracture (rare) • Osteoporosis diagnosis in children requires clinical evidence of bone fragility (e.g., fracture with minimal trauma) and cannot rely solely on measured low BMD.	<u>Low BMD:</u> • 7% of a U.S. cohort had a BMD z score less than or equal to -2.0 (87% treated with ART). • 24% to 32% of Thai and Brazilian adolescents had a BMD z score less than or equal to -2.0 (92% to 100% treated with ART).	Longer duration of HIV infection Greater severity of HIV disease Growth delay, pubertal delay Low BMI Lipodystrophy Non-black race Smoking Prolonged systemic corticosteroid use Medroxyprogesterone use Limited weight-bearing exercise	<u>Prevention:</u> • Ensure sufficient calcium and vitamin D intake. • Encourage weight-bearing exercise. • Minimize modifiable risk factors (e.g., smoking, low BMI, steroid use). <u>Monitoring:</u> • Assess nutritional intake (calcium, vitamin D, and total calories). • Consider obtaining serum 25-OH-vitamin D level. ^a • Obtain DXA. ^b	Ensure sufficient calcium intake and vitamin D sufficiency. Encourage weight-bearing exercise. Reduce modifiable risk factors (e.g., smoking, low BMI, use of steroids, use of medroxyprogesterone). Role of bisphosphonates not established in children Consider change in ARV regimen.

^a Some experts would periodically measure 25-OH-vitamin D, especially in HIV-infected urban youth, because in that population, the prevalence of vitamin D insufficiency is high.

^b Until more data are available about the long-term effects of TDF on bone mineral acquisition in childhood, some experts would obtain a DXA at baseline and every 6 to 12 months for prepubertal children and children in early puberty who are initiating treatment with TDF. DXA could also be considered in adolescent women on TDF and medroxyprogesterone and in children with indications not uniquely related to HIV infection (such as cerebral palsy).

Key to Acronyms: ART = antiretroviral therapy; ARV = antiretroviral; BMD = bone mineral density; BMI = body mass index; DXA = dual-energy x-ray absorptiometry; LPV/r = lopinavir/ritonavir; PI = protease inhibitor; TDF = tenofovir disoproxil fumarate

References

Osteopenia and Osteoporosis

- McComsey GA, Tebas P, Shane E, et al. Bone disease in HIV infection: a practical review and recommendations for HIV care providers. *Clin Infect Dis*. 2010;51(8):937-946. Available at <http://www.ncbi.nlm.nih.gov/pubmed/20839968>.
- Mora S, Zamproni I, Beccio S, Bianchi R, Giacomet V, Vigano A. Longitudinal changes of bone mineral density and metabolism in antiretroviral-treated human immunodeficiency virus-infected children. *The Journal of clinical endocrinology and metabolism*. 2004;89(1):24-28. Available at

<http://www.ncbi.nlm.nih.gov/pubmed/14715822>.

3. Gafni RI, Hazra R, Reynolds JC, et al. Tenofovir disoproxil fumarate and an optimized background regimen of antiretroviral agents as salvage therapy: impact on bone mineral density in HIV-infected children. *Pediatrics*. 2006;118(3):e711-718. Available at <http://www.ncbi.nlm.nih.gov/pubmed/16923923>.
4. Purdy JB, Gafni RI, Reynolds JC, Zeichner S, Hazra R. Decreased bone mineral density with off-label use of tenofovir in children and adolescents infected with human immunodeficiency virus. *J Pediatr*. 2008;152(4):582-584. Available at <http://www.ncbi.nlm.nih.gov/pubmed/18346519>.
5. Jacobson DL, Lindsey JC, Gordon CM, et al. Total body and spinal bone mineral density across Tanner stage in perinatally HIV-infected and uninfected children and youth in PACTG 1045. *AIDS*. 2010;24(5):687-696. Available at <http://www.ncbi.nlm.nih.gov/pubmed/20168204>.
6. Jacobson DL, Spiegelman D, Duggan C, et al. Predictors of bone mineral density in human immunodeficiency virus-1 infected children. *Journal of pediatric gastroenterology and nutrition*. 2005;41(3):339-346. Available at <http://www.ncbi.nlm.nih.gov/pubmed/16131991>.
7. Kalkwarf HJ, Zemel BS, Gilsanz V, et al. The bone mineral density in childhood study: bone mineral content and density according to age, sex, and race. *The Journal of clinical endocrinology and metabolism*. 2007;92(6):2087-2099. Available at <http://www.ncbi.nlm.nih.gov/pubmed/17311856>.
8. Bachrach LK, Sills IN, Section on E. Clinical report-bone densitometry in children and adolescents. *Pediatrics*. 2011;127(1):189-194. Available at <http://www.ncbi.nlm.nih.gov/pubmed/21187316>.
9. Lima LR, Silva RC, Giuliano Ide C, Sakuno T, Brincas SM, Carvalho AP. Bone mass in children and adolescents infected with human immunodeficiency virus. *Jornal de pediatria*. 2013;89(1):91-99. Available at <http://www.ncbi.nlm.nih.gov/pubmed/23544816>.
10. Puthanakit T, Saksawad R, Bunupuradah T, et al. Prevalence and risk factors of low bone mineral density among perinatally HIV-infected Thai adolescents receiving antiretroviral therapy. *J Acquir Immune Defic Syndr*. 2012;61(4):477-483. Available at <http://www.ncbi.nlm.nih.gov/pubmed/22918157>.
11. Siberry GK, Li H, Jacobson D, Pediatric ACTGCS. Fracture risk by HIV infection status in perinatally HIV-exposed children. *AIDS Res Hum Retroviruses*. 2012;28(3):247-250. Available at <http://www.ncbi.nlm.nih.gov/pubmed/22471877>.
12. DiMeglio LA, Wang J, Siberry GK, et al. Bone mineral density in children and adolescents with perinatal HIV infection. *AIDS*. 2013;27(2):211-220. Available at <http://www.ncbi.nlm.nih.gov/pubmed/23032412>.
13. Bunders MJ, Frinking O, Scherpbier HJ, et al. Bone mineral density increases in HIV-infected children treated with long-term combination antiretroviral therapy. *Clin Infect Dis*. 2013;56(4):583-586. Available at <http://www.ncbi.nlm.nih.gov/pubmed/23097583>.
14. Huang JS, Hughes MD, Riddler SA, Haubrich RH, Aids Clinical Trials Group AST. Bone mineral density effects of randomized regimen and nucleoside reverse transcriptase inhibitor selection from ACTG A5142. *HIV Clin Trials*. 2013;14(5):224-234. Available at <http://www.ncbi.nlm.nih.gov/pubmed/24144899>.
15. Puthanakit T, Siberry GK. Bone health in children and adolescents with perinatal HIV infection. *J Int AIDS Soc*. 2013;16:18575. Available at <http://www.ncbi.nlm.nih.gov/pubmed/23782476>.
16. Aurpibul L, Cressey TR, Sricharoenchai S, et al. Efficacy, safety and pharmacokinetics of tenofovir disoproxil fumarate in virologic-suppressed HIV-infected children using weight-band dosing. *Pediatr Infect Dis J*. 2015;34(4):392-397. Available at <http://www.ncbi.nlm.nih.gov/pubmed/25760566>.
17. Overton ET, Chan ES, Brown TT, et al. Vitamin D and calcium attenuate bone loss with antiretroviral therapy initiation: a randomized trial. *Ann Intern Med*. 2015;162(12):815-824. Available at <http://www.ncbi.nlm.nih.gov/pubmed/26075752>.
18. Ross AC. The 2011 report on dietary reference intakes for calcium and vitamin D. *Public Health Nutr*. 2011;14(5):938-939. Available at <http://www.ncbi.nlm.nih.gov/pubmed/21492489>.

Table 12k. Antiretroviral-Therapy-Associated Adverse Effects and Management Recommendations—Peripheral Nervous System Toxicity (Last updated March 1, 2016; last reviewed March 1, 2016)

Adverse Effects	Associated ARVs	Onset/Clinical Manifestations	Estimated Frequency ^a	Risk Factors	Prevention/Monitoring	Management
ARV Toxic Neuropathy^b	d4T, ddI PIs	<p><u>Onset:</u></p> <ul style="list-style-type: none"> Variable; weeks to months following NRTI initiation. <p><u>Presentation:</u></p> <ul style="list-style-type: none"> Decreased sensation Aching, burning, painful numbness Hyperalgesia (lowered pain threshold) Allodynia (non noxious stimuli cause pain) Decreased or absent ankle reflexes <p><u>Distribution:</u></p> <ul style="list-style-type: none"> Bilateral soles of feet, ascending to legs and fingertips 	<p><u>HIV-Infected Children:</u></p> <ul style="list-style-type: none"> 1.13% prevalence (baseline 2001); incidence 0.23 per 100 person-years (2001–2006) in a U.S. cohort. <1% discontinued d4T because of neuropathy in 3 large African cohorts (aged 1 month–18 years; median follow-up 1.8–3.2 years). 42 out of 174 (24%) in a South African cohort were diagnosed with peripheral neuropathy. 86% were taking d4T, and use of ddI was an additional risk factor. 4/40 (10%) Indian children taking d4T had abnormal nerve conduction tests. <p><u>HIV-Infected Adults on d4T:</u></p> <ul style="list-style-type: none"> Prevalence up to 57% Incidence rates of 6.4–12.1 per 100 person-years 	<p><u>HIV-Infected Adults:</u></p> <ul style="list-style-type: none"> Preexisting neuropathy (e.g., diabetes, alcohol abuse, vitamin B-12 deficiency) Elevated triglyceride levels Older age Poor nutrition More advanced HIV disease Concomitant use of other neurotoxic agents (e.g., INH) Some mitochondrial DNA haplogroups may have increased risk. 	<p>Limit use of d4T and ddI.</p> <p>As part of routine care, monitor for symptoms and signs of peripheral neuropathy.</p>	<p>Discontinue offending agent.</p> <p>Persistent pain can be difficult to treat; topical capsaicin 8% may be helpful.</p> <p>Consider referral to a neurologist.</p> <p>Data Are Insufficient to Allow the Panel to Recommend Use of Any of the Following Modalities in Children:</p> <ul style="list-style-type: none"> Tricyclic antidepressants Gabapentin Pregabalin Mexiletine Lamotrigine Acupuncture or other complementary approaches

^a Peripheral neuropathy may be underreported in children because symptoms are difficult to evaluate in young children.

^b HIV infection itself may cause a distal sensory neuropathy that is phenotypically identical to ARV toxic neuropathy.

Key to Acronyms: ARV = antiretroviral; d4T = stavudine; ddI = didanosine; INH = isoniazid; NRTI = nucleoside reverse transcriptase inhibitor; PI = protease inhibitor

References

1. Nachman SA, Chernoff M, Gona P, et al. Incidence of noninfectious conditions in perinatally HIV-infected children and adolescents in the HAART era. *Arch Pediatr Adolesc Med.* 2009;163(2):164-171. Available at <http://www.ncbi.nlm.nih.gov/pubmed/19188649>.
2. Buck WC, Kabue MM, Kazembe PN, Kline MW. Discontinuation of standard first-line antiretroviral therapy in a cohort of 1434 Malawian children. *J Int AIDS Soc.* 2010;13:31. Available at <http://www.ncbi.nlm.nih.gov/pubmed/20691049>.
3. Ances BM, Vaida F, Rosario D, et al. Role of metabolic syndrome components in HIV-associated sensory neuropathy. *AIDS.* 2009;23(17):2317-2322. Available at <http://www.ncbi.nlm.nih.gov/pubmed/19823068>.
4. Banerjee S, McCutchan JA, Ances BM, et al. Hypertriglyceridemia in combination antiretroviral-treated HIV-positive individuals: potential impact on HIV sensory polyneuropathy. *AIDS.* 2011;25(2):F1-6. Available at <http://www.ncbi.nlm.nih.gov/pubmed/21150557>.
5. Canter JA, Robbins GK, Selph D, et al. African mitochondrial DNA subhaplogroups and peripheral neuropathy during antiretroviral therapy. *J Infect Dis.* 2010;201(11):1703-1707. Available at <http://www.ncbi.nlm.nih.gov/pubmed/20402593>.
6. McCormack PL. Capsaicin dermal patch: in non-diabetic peripheral neuropathic pain. *Drugs.* 2010;70(14):1831-1842. Available at <http://www.ncbi.nlm.nih.gov/pubmed/20836576>.
7. Phillips TJ, Cherry CL, Cox S, Marshall SJ, Rice AS. Pharmacological treatment of painful HIV-associated sensory neuropathy: a systematic review and meta-analysis of randomised controlled trials. *PLoS One.* 2010;5(12):e14433. Available at <http://www.ncbi.nlm.nih.gov/pubmed/21203440>.
8. Menezes CN, Maskew M, Sanne I, Crowther NJ, Raal FJ. A longitudinal study of stavudine-associated toxicities in a large cohort of South African HIV infected subjects. *BMC Infect Dis.* 2011;11:244. Available at <http://www.ncbi.nlm.nih.gov/pubmed/21923929>.
9. Wadley AL, Cherry CL, Price P, Kamerman PR. HIV neuropathy risk factors and symptom characterization in stavudine-exposed South Africans. *Journal of pain and symptom management.* 2011;41(4):700-706. Available at <http://www.ncbi.nlm.nih.gov/pubmed/21145196>.
10. Tukei VJ, Asiimwe A, Maganda A, et al. Safety and tolerability of antiretroviral therapy among HIV-infected children and adolescents in Uganda. *J Acquir Immune Defic Syndr.* 2012;59(3):274-280. Available at <http://www.ncbi.nlm.nih.gov/pubmed/22126740>.
11. Webster LR, Peppin JF, Murphy FT, Tobias JK, Vanhove GF. Tolerability of NGX-4010, a capsaicin 8% patch, in conjunction with three topical anesthetic formulations for the treatment of neuropathic pain. *Journal of pain research.* 2012;5:7-13. Available at <http://www.ncbi.nlm.nih.gov/pubmed/22328830>.
12. Phan V, Thai S, Choun K, Lynen L, van Griensven J. Incidence of treatment-limiting toxicity with stavudine-based antiretroviral therapy in Cambodia: a retrospective cohort study. *PLoS One.* 2012;7(1):e30647. Available at <http://www.ncbi.nlm.nih.gov/pubmed/22303447>.
13. Palmer M, Chersich M, Moultrie H, Kuhn L, Fairlie L, Meyers T. Frequency of stavudine substitution due to toxicity in children receiving antiretroviral treatment in sub-Saharan Africa. *AIDS.* 2013;27(5):781-785. Available at <http://www.ncbi.nlm.nih.gov/pubmed/23169331>.
14. Peters RP, Van Ramshorst MS, Struthers HE, McIntyre JA. Clinical assessment of peripheral neuropathy in HIV-infected children on antiretroviral therapy in rural South Africa. *Eur J Pediatr.* 2014;173(9):1245-1248. Available at <http://www.ncbi.nlm.nih.gov/pubmed/24691679>.
15. Kaku M, Simpson DM. HIV neuropathy. *Curr Opin HIV AIDS.* 2014;9(6):521-526. Available at <http://www.ncbi.nlm.nih.gov/pubmed/25275705>.
16. Sankhyan N, Lodha R, Sharma S, et al. Peripheral neuropathy in children on stavudine therapy. *Indian journal of pediatrics.* 2015;82(2):136-139. Available at <http://www.ncbi.nlm.nih.gov/pubmed/24874810>.
17. Chen H, Clifford DB, Deng L, et al. Peripheral neuropathy in ART-experienced patients: prevalence and risk factors. *J Neurovirol.* 2013;19(6):557-564. Available at <http://www.ncbi.nlm.nih.gov/pubmed/24297499>.

Table 12I. Antiretroviral-Therapy-Associated Adverse Effects and Management Recommendations—Rash and Hypersensitivity Reactions (Last updated March 5, 2015; last reviewed March 5, 2015) (page 1 of 4)

Adverse Effects	Associated ARVs	Onset/Clinical Manifestations	Estimated Frequency	Risk Factors	Prevention/Monitoring	Management
Rash	Any ARV can cause rash.	<p><u>Onset:</u></p> <ul style="list-style-type: none"> First few days to weeks after starting therapy <p><u>Presentation:</u></p> <ul style="list-style-type: none"> Most rashes are mild-to-moderate, diffuse maculopapular eruptions. <p>Note: Some rashes are the initial manifestation of systemic hypersensitivity (see Systemic HSR, SJS/TEN/EM Major).</p>	<p><u>Common (>10% Adults and/or Children):</u></p> <ul style="list-style-type: none"> NVP, EFV, ETR, FPV, FTC <p><u>Less Common (5% to 10%):</u></p> <ul style="list-style-type: none"> ABC, DRV, TPV, TDF <p><u>Unusual (2% to 4%):</u></p> <ul style="list-style-type: none"> LPV/r, RAL, MVC, RPV 	<ul style="list-style-type: none"> Sulfonamide allergy is a risk factor for rash with PIs containing a sulfonamide moiety (FPV, DRV, and TPV). Possible association of polymorphisms in CYP2B6 and multiple HLA loci with rash with NVP. 	<p><u>When Starting NVP or Restarting After Interruptions >14 Days:</u></p> <ul style="list-style-type: none"> Once-daily dosing (50% of total daily dose) for 2 weeks, then escalation to target dose with twice-daily dosing is associated with fewer rashes.^a Avoid the use of systemic corticosteroids during NVP dose escalation. Assess patient for rash severity, mucosal involvement, and other signs of systemic reaction. Consider concomitant medications and illnesses that cause rash. 	<p><u>Mild-To-Moderate Maculopapular Rash Without Systemic or Mucosal Involvement:</u></p> <ul style="list-style-type: none"> Most will resolve without intervention; ARVs can be continued while monitoring.^a Antihistamines may provide some relief. <p><u>Severe Rash (e.g., Blisters, Bullae, Ulcers, Skin Necrosis) and/or Rash Accompanied by Systemic Symptoms (e.g., Fever, Arthralgias, Edema) and/or Rash Accompanied By Mucous Membrane Involvement (e.g., Conjunctivitis):</u></p> <ul style="list-style-type: none"> Manage as SJS/TEN/EM major (see below). <p><u>Rash in Patients Receiving NVP:</u></p> <ul style="list-style-type: none"> Given elevated risk of HSR, measure hepatic transaminases. If hepatic transaminases are elevated, NVP should be discontinued and not restarted (see HSR-NVP).
	ENF	<p><u>Onset:</u></p> <ul style="list-style-type: none"> First few days to weeks after starting therapy <p><u>Presentation:</u></p> <ul style="list-style-type: none"> Local injection site reactions with pain, erythema, induration, nodules and cysts, pruritus, ecchymosis. Often multiple reactions at the same time. 	<p><u>Adults and Children:</u></p> <ul style="list-style-type: none"> >90% 	Unknown	<ul style="list-style-type: none"> Routinely assess patient for local reactions. Rotate injection sites. Massage area after injection. 	<ul style="list-style-type: none"> Continue the agent as tolerated by the patient. Adjust injection technique. Rotate injection sites.

Table 12I. Antiretroviral-Therapy-Associated Adverse Effects and Management Recommendations—Rash and Hypersensitivity Reactions (Last updated March 5, 2015; last reviewed March 5, 2015) (page 2 of 4)

Adverse Effects	Associated ARVs	Onset/Clinical Manifestations	Estimated Frequency	Risk Factors	Prevention/Monitoring	Management
SJS/TEN/EM Major	Many ARVs, especially NNRTIs (see frequency column)	<p><u>Onset:</u></p> <ul style="list-style-type: none"> First few days to weeks after initiating therapy <p><u>Presentation:</u></p> <ul style="list-style-type: none"> Initial rash may be mild, but often becomes painful, evolving to blister/bulla formation with necrosis in severe cases. Usually involves mucous membrane ulceration and/or conjunctivitis. Systemic symptoms may include fever, tachycardia, malaise, myalgia, and arthralgia. 	<p><u>Infrequent:</u></p> <ul style="list-style-type: none"> NVP (0.3%), EFV (0.1%), ETR (<0.1%) <p><u>Case Reports:</u></p> <ul style="list-style-type: none"> FPV, ABC, DRV, ZDV, ddI, IDV, LPV/r, ATV, RAL 	<p><u>Adults:</u></p> <ul style="list-style-type: none"> Female gender Race/ethnicity (black, Asian, Hispanic) 	<p>To Lower the Risk of Reactions to NVP when Starting or Restarting after Interruptions >14 Days:</p> <ul style="list-style-type: none"> Utilize once-daily dosing (50% of total daily dose) for 2 weeks, then escalate to target dose with twice-daily dosing, which is associated with fewer rashes.^a Counsel families to report symptoms as soon as they appear. 	<ul style="list-style-type: none"> Discontinue all ARVs and other possible causative agents such as cotrimoxazole. Provide intensive supportive care, IV hydration, aggressive wound care, pain management, antipyretics, parenteral nutrition, and antibiotics as needed in case of superinfection. Corticosteroids and/or IVIG are sometimes used, but use of each is controversial. Do not reintroduce the offending medication. In case of SJS/TEN/EM major with one NNRTI, many experts would avoid use of other NNRTIs.
<p>Systemic HSR</p> <p>With or without skin involvement and excluding SJS/TEN</p>	ABC	<p><u>Onset</u></p> <p><i>With First Use:</i></p> <ul style="list-style-type: none"> Within first 6 weeks <p><i>With Re-Introduction:</i></p> <ul style="list-style-type: none"> Within hours <p><u>Presentation:</u></p> <ul style="list-style-type: none"> Symptoms include high fever, diffuse skin rash, malaise, nausea, headache, myalgia, arthralgia, diarrhea, vomiting, abdominal pain, pharyngitis, respiratory symptoms (e.g., dyspnea). Symptoms worsen to include hypotension and vascular collapse with continuation. With re-challenge, symptoms can mimic anaphylaxis. 	2.3% to 9% (varies by racial/ethnic group).	<ul style="list-style-type: none"> HLA-B*5701 (HSR very uncommon in people who are HLA-B*5701-negative); also HLA-DR7, HLA-DQ3. HSR risk is higher in those of white race compared to those of black or East Asian race. 	<ul style="list-style-type: none"> Screening for HLA-B*5701. ABC should not be prescribed if HLA-B*5701 is positive. The medical record should clearly indicate that ABC is contraindicated. When starting ABC, counsel patients and families about the signs and symptoms of HSR to ensure prompt reporting of reactions. 	<ul style="list-style-type: none"> Discontinue ARVs and investigate for other causes of the symptoms (e.g., a concurrent viral illness). Treat symptoms as necessary. Most symptoms resolve within 48 hours after discontinuation of ABC. Do not re-challenge with ABC even if the patient is HLA-B*5701-negative.

Table 12I. Antiretroviral-Therapy-Associated Adverse Effects and Management Recommendations—Rash and Hypersensitivity Reactions (Last updated March 5, 2015; last reviewed March 5, 2015) (page 3 of 4)

Adverse Effects	Associated ARVs	Onset/Clinical Manifestations	Estimated Frequency	Risk Factors	Prevention/Monitoring	Management
Systemic HSR With or without skin involvement and excluding SJS/TEN	NVP	<u>Onset:</u> <ul style="list-style-type: none"> Most frequent in the first few weeks of therapy but can occur through 18 weeks. <u>Presentation:</u> <ul style="list-style-type: none"> Flu-like symptoms (including nausea, vomiting, myalgia, fatigue, fever, abdominal pain, jaundice) with or without skin rash that may progress to hepatic failure with encephalopathy. DRESS syndrome has also been described. 	4% (2.5% to 11%)	<u>Adults:</u> <ul style="list-style-type: none"> Treatment-naïve with higher CD4 count (>250 cells/mm³ in women; >400 cells/mm³ in men). Female gender (risk is 3-fold higher in females compared with males). <u>Children:</u> <ul style="list-style-type: none"> NVP hepatotoxicity and HSR are less common in pre-pubertal children than in adults. The PREDICT Study showed a 2.65 times higher risk of overall NVP toxicity (rash, hepatotoxicity, hypersensitivity) in children with CD4 $\geq 15\%$ compared to children with CD4 $<15\%$. 	<u>When Starting NVP or Restarting After Interruptions ≥ 14 Days:</u> <ul style="list-style-type: none"> Two-week lead-in period with once-daily dosing then dose escalation to twice daily as recommended may reduce risk of reaction.^a Counsel families about signs and symptoms of HSR to ensure prompt reporting of reactions. Obtain AST and ALT in patients with rash. Obtain AST and ALT at baseline, before dose escalation, 2 weeks post-dose escalation, and thereafter at 3-month intervals. Avoid NVP use in women with CD4 counts >250 cells/mm³ and in men with CD4 counts >400 cells/mm³ unless benefits outweigh risks. Do not use NVP in PEP. 	<ul style="list-style-type: none"> Discontinue ARVs. Consider other causes for hepatitis and discontinue all hepatotoxic medications. Provide supportive care as indicated and monitor patient closely. Do not re-introduce NVP. The safety of other NNRTIs is unknown following symptomatic hepatitis due to NVP, and many experts would avoid the NNRTI drug class when restarting treatment.
	ENF, ETR	<u>Onset:</u> <ul style="list-style-type: none"> Any time during therapy. <u>Presentation:</u> <ul style="list-style-type: none"> Symptoms may include rash, constitutional findings, and sometimes organ dysfunction including hepatic failure. 	Rare	Unknown	Evaluate for hypersensitivity if the patient is symptomatic.	Discontinue ARVs. Re-challenge with ENF or ETR is not recommended.

Table 12I. Antiretroviral-Therapy-Associated Adverse Effects and Management Recommendations—Rash and Hypersensitivity Reactions (Last updated March 5, 2015; last reviewed March 5, 2015) (page 4 of 4)

Adverse Effects	Associated ARVs	Onset/Clinical Manifestations	Estimated Frequency	Risk Factors	Prevention/Monitoring	Management
Systemic HSR With or without skin involvement and excluding SJS/TEN	RAL	DRESS syndrome	Rare	Unknown	Evaluate for hypersensitivity if the patient is symptomatic.	Discontinue all ARVs. Re-challenge with RAL is not recommended.
	MVC	Rash preceding hepatotoxicity	Rare	Unknown	Obtain AST and ALT in patients with rash or other symptoms of hypersensitivity.	Discontinue all ARVs. Re-challenge with MVC is not recommended.

^a The prescribing information for NVP states that patients experiencing rash during the 14-day lead-in period should not have the NVP dose increased until the rash has resolved. However, prolonging the lead-in phase beyond 14 days may increase risk of NVP resistance because of sub-therapeutic drug levels. Management of children who have persistent mild or moderate rash after the lead-in period should be individualized and consultation with an expert in HIV care should be obtained. **NVP should be stopped and not restarted** if the rash is severe or is worsening or progressing.

Key to Acronyms: ABC = abacavir; ALT = alanine transaminase; ARV = antiretroviral; AST = aspartate aminotransferase; ATV = atazanavir; CD4 = CD4 T lymphocyte cell; ddI = didanosine; DRESS = drug rash with eosinophilia and systemic symptoms; DRV = darunavir; EFV = efavirenz; EM = erythema multiforme; ENF = enfuvirtide; ETR = etravirine; FPV = fosamprenavir; FTC = emtricitabine; HSR = hypersensitivity reaction; IDV = indinavir; IV = intravenous; IVIG = intravenous immune globulin; LPV/r = lopinavir/ritonavir; MVC = maraviroc; NNRTI = non-nucleoside reverse transcriptase inhibitor; NVP = nevirapine; PEP = post-exposure prophylaxis; PI = protease inhibitor; RAL = raltegravir; RPV = rilpivirine; SJS = Stevens-Johnson syndrome; TDF = tenofovir disoproxil fumarate; TEN = toxic epidermal necrolysis; TPV = tipranavir; ZDV = zidovudine

References

1. Borras-Blasco J, Navarro-Ruiz A, Borras C, Castera E. Adverse cutaneous reactions associated with the newest antiretroviral drugs in patients with human immunodeficiency virus infection. *J Antimicrob Chemother*. 2008;62(5):879-888. Available at <http://www.ncbi.nlm.nih.gov/pubmed/18653488>.
2. Davis CM, Shearer WT. Diagnosis and management of HIV drug hypersensitivity. *J Allergy Clin Immunol*. 2008;121(4):826-832 e825. Available at <http://www.ncbi.nlm.nih.gov/pubmed/18190954>.
3. Kea C, Puthanakit T, Apornpong T, et al. Incidence and risk factors for nevirapine related toxicities among HIV-infected Asian children randomized to starting ART at different CD4%. Abstract MOPE240. Presented at: 6th International AIDS Society Conference on HIV Pathogenesis and Treatment and Prevention. 2011. Rome, Italy.
4. Mallal S, Nolan D, Witt C, et al. Association between presence of HLA-B*5701, HLA-DR7, and HLA-DQ3 and hypersensitivity to HIV-1 reverse-transcriptase inhibitor abacavir. *Lancet*. 2002;359(9308):727-732. Available at <http://www.ncbi.nlm.nih.gov/pubmed/11888582>.
5. Mallal S, Phillips E, Carosi G, et al. HLA-B*5701 screening for hypersensitivity to abacavir. *N Engl J Med*. 2008;358(6):568-579. Available at <http://www.ncbi.nlm.nih.gov/pubmed/18256392>.

6. Mirochnick M, Clarke DF, Dorenbaum A. Nevirapine: pharmacokinetic considerations in children and pregnant women. *Clinical pharmacokinetics*. 2000;39(4):281-293. Available at <http://www.ncbi.nlm.nih.gov/pubmed/11069214>.
7. Puthanakit T, Bunupuradah T, Kosalaraksa P, et al. Prevalence of human leukocyte antigen-B*5701 among HIV-infected children in Thailand and Cambodia: implications for abacavir use. *Pediatr Infect Dis J*. 2013;32(3):252-253. Available at <http://www.ncbi.nlm.nih.gov/pubmed/22986704>.
8. Stern JO, Robinson PA, Love J, Lanes S, Imperiale MS, Mayers DL. A comprehensive hepatic safety analysis of nevirapine in different populations of HIV infected patients. *J Acquir Immune Defic Syndr*. 2003;34 Suppl 1(Suppl 1):S21-33. Available at <http://www.ncbi.nlm.nih.gov/pubmed/14562855>.
9. Shubber Z, Calmy A, Andrieux-Meyer I, et al. Adverse events associated with nevirapine and efavirenz-based first-line antiretroviral therapy: a systematic review and meta-analysis. *AIDS*. 2013;27(9):1403-1412. Available at <http://www.ncbi.nlm.nih.gov/pubmed/23343913>.
10. Tas S, Simonart T. Management of drug rash with eosinophilia and systemic symptoms (DRESS syndrome): an update. *Dermatology*. 2003;206(4):353-356. Available at <http://www.ncbi.nlm.nih.gov/pubmed/12771485>.
11. Trottier B, Walmsley S, Reynes J, et al. Safety of enfuvirtide in combination with an optimized background of antiretrovirals in treatment-experienced HIV-1-infected adults over 48 weeks. *J Acquir Immune Defic Syndr*. 2005;40(4):413-421. Available at <http://www.ncbi.nlm.nih.gov/pubmed/16280695>.
12. Vitezica ZG, Milpied B, Lonjou C, et al. HLA-DRB1*01 associated with cutaneous hypersensitivity induced by nevirapine and efavirenz. *AIDS*. 2008;22(4):540-541. Available at <http://www.ncbi.nlm.nih.gov/pubmed/18301070>.
13. Yuan J, Guo S, Hall D, et al. Toxicogenomics of nevirapine-associated cutaneous and hepatic adverse events among populations of African, Asian, and European descent. *AIDS*. 2011;25(10):1271-1280. Available at <http://www.ncbi.nlm.nih.gov/pubmed/21505298>.
14. Dziuban EJ, Hughey AB, Stewart DA, et al. Stevens-Johnson syndrome and HIV in children in Swaziland. *Pediatr Infect Dis J*. 2013;32(12):1354-1358. Available at <http://www.ncbi.nlm.nih.gov/pubmed/23743542>.
15. Rutstein RM, Samson P, Fenton T, et al; for the PACTG 1020A Study Team. Long-term safety and efficacy of atazanavir-based therapy in HIV-infected infants, children and adolescents: the pediatric aids clinical trials group protocol 1020A. *Pediatr Infect Dis J*. 2015;34:162-167. Available at <http://www.ncbi.nlm.nih.gov/pubmed/25232777>.